

PROFINITE GROUPS
WITH A CYCLOTOMIC p -ORIENTATION

TO THE MEMORY OF VLADIMIR VOEVODSKY

CLAUDIO QUADRELLI AND THOMAS S. WEIGEL

Received: November 6, 2018

Revised: September 12, 2020

Communicated by Otmar Venjakob

ABSTRACT. Let p be a prime. A continuous representation $\theta: G \rightarrow \mathrm{GL}_1(\mathbb{Z}_p)$ of a profinite group G is called a cyclotomic p -orientation if for all open subgroups $U \subseteq G$ and for all $k, n \geq 1$ the natural maps $H^k(U, \mathbb{Z}_p(k)/p^n) \rightarrow H^k(U, \mathbb{Z}_p(k)/p)$ are surjective. Here $\mathbb{Z}_p(k)$ denotes the \mathbb{Z}_p -module of rank 1 with U -action induced by $\theta|_U^k$. By the Rost-Voevodsky theorem, the cyclotomic character of the absolute Galois group $G_{\mathbb{K}}$ of a field \mathbb{K} is, indeed, a cyclotomic p -orientation of $G_{\mathbb{K}}$. We study profinite groups with a cyclotomic p -orientation. In particular, we show that cyclotomicity is preserved by several operations on profinite groups, and that Bloch-Kato pro- p groups with a cyclotomic p -orientation satisfy a strong form of Tits' alternative and decompose as semi-direct product over a canonical abelian closed normal subgroup.

2020 Mathematics Subject Classification: Primary 12G05; Secondary 20E18, 12F10

Keywords and Phrases: Absolute Galois groups, Rost-Voevodsky theorem, elementary type conjecture

1 INTRODUCTION

For a prime p let \mathbb{Z}_p denote the ring of p -adic integers. For a profinite group G , we call a continuous representation $\theta: G \rightarrow \mathbb{Z}_p^\times = \mathrm{GL}_1(\mathbb{Z}_p)$ a p -orientation of G and call the couple (G, θ) a p -oriented profinite group. Given a p -oriented

profinite group (G, θ) , for $k \in \mathbb{Z}$ let $\mathbb{Z}_p(k)$ denote the left $\mathbb{Z}_p[[G]]$ -module induced by θ^k , namely, $\mathbb{Z}_p(k)$ is equal to the additive group \mathbb{Z}_p and the left G -action is given by

$$g \cdot z = \theta(g)^k \cdot z, \quad g \in G, z \in \mathbb{Z}_p(k). \tag{1.1}$$

Vice-versa, if M is a topological left $\mathbb{Z}_p[[G]]$ -module which as an abelian pro- p group is isomorphic to \mathbb{Z}_p , then there exists a unique p -orientation $\theta: G \rightarrow \mathbb{Z}_p^\times$ such that $M \simeq \mathbb{Z}_p(1)$.

The $\mathbb{Z}_p[[G]]$ -module $\mathbb{Z}_p(1)$ and the representation $\theta: G \rightarrow \mathbb{Z}_p^\times$ are said to be k -cyclotomic, for $k \geq 1$, if for every open subgroup U of G and every $n \geq 1$ the natural maps

$$H^k(U, \mathbb{Z}_p(k)/p^n) \longrightarrow H^k(U, \mathbb{Z}_p(k)/p), \tag{1.2}$$

induced by the epimorphism of $\mathbb{Z}_p[[U]]$ -modules $\mathbb{Z}_p(k)/p^n \rightarrow \mathbb{Z}_p(k)/p$, are surjective. If $\mathbb{Z}_p(1)$ (respectively θ) is k -cyclotomic for every $k \geq 1$, then it is called simply a cyclotomic $\mathbb{Z}_p[[G]]$ -module (resp., cyclotomic p -orientation). Note that $\mathbb{Z}_p(1)$ is k -cyclotomic if, and only if, $H_{\text{cts}}^{k+1}(U, \mathbb{Z}_p(k))$ is a torsion free \mathbb{Z}_p -module for every open subgroup $U \subseteq G$ — here H_{cts}^* denotes continuous cochain cohomology as introduced by J. Tate in [34] (see § 2.1).

Cyclotomic modules of profinite groups have been introduced and studied by C. De Clercq and M. Florence in [5]. Previously J.P. Labute, in [16], considered surjectivity of (1.2) in the case $k = 1$ and $U = G$ — note that demanding surjectivity for $U = G$ only is much weaker than demanding it for every open subgroup $U \subseteq G$, and this is what makes the definition of cyclotomic modules truly new.

Let \mathbb{K} be a field, and let $\bar{\mathbb{K}}/\mathbb{K}$ be a separable closure of \mathbb{K} . If $\text{char}(\mathbb{K}) \neq p$, the absolute Galois group $G_{\mathbb{K}} = \text{Gal}(\bar{\mathbb{K}}/\mathbb{K})$ of \mathbb{K} comes equipped with a canonical p -orientation

$$\theta_{\mathbb{K},p}: G_{\mathbb{K}} \longrightarrow \text{Aut}(\mu_{p^\infty}(\bar{\mathbb{K}})) \simeq \mathbb{Z}_p^\times, \tag{1.3}$$

where $\mu_{p^\infty}(\bar{\mathbb{K}}) \subseteq \bar{\mathbb{K}}^\times$ denotes the subgroup of roots of unity of $\bar{\mathbb{K}}$ of p -power order. If $p = \text{char}(\mathbb{K})$, we put $\theta_{\mathbb{K},p} = \mathbf{1}_{G_{\mathbb{K}}}$, the function which is constantly 1 on $G_{\mathbb{K}}$. The following result (cf. [5, Prop. 14.19]) is a consequence of the positive solution of the Bloch-Kato Conjecture given by M. Rost and V. Voevodsky with the ‘‘C. Weibel patch’’ (cf. [29, 36, 40]), which from now on we will refer to as the Rost-Voevodsky Theorem.

THEOREM 1.1. *Let \mathbb{K} be a field, and let p be prime number. The canonical p -orientation $\theta_{\mathbb{K},p}: G_{\mathbb{K}} \rightarrow \mathbb{Z}_p^\times$ is cyclotomic.*

Theorem 1.1 provides a fundamental class of examples of profinite groups endowed with a cyclotomic p -orientation. Bearing in mind the exotic character of absolute Galois groups, it also provides a strong motivation to the study of cyclotomically p -oriented profinite groups — which is the main purpose of this manuscript. In fact, one may recover several Galois-theoretic statements already for profinite groups with a 1-cyclotomic p -orientation — e.g., the only finite group endowed with a 1-cyclotomic p -orientation is the finite group C_2

of order 2, with non-constant 2-orientation $\theta: C_2 \rightarrow \{\pm 1\}$ (cf. [11, Ex. 3.5]), and this implies the Artin-Schreier obstruction for absolute Galois groups. In their paper, De Clercq and Florence formulated the ‘‘Smoothness Conjecture’’, which can be restated in this context as follows: for a p -oriented profinite group, 1-cyclotomicity implies k -cyclotomicity for all $k \geq 1$ (cf. [5, Conj. 14.25]). A p -oriented profinite group (G, θ) is said to be *Bloch-Kato* if the \mathbb{F}_p -algebra

$$H^\bullet(U, \widehat{\theta}|_U) = \prod_{k \geq 0} H^k(U, \mathbb{F}_p(k)), \quad (1.4)$$

where $\mathbb{F}_p(k) = \mathbb{Z}_p(k)/p$, with product given by cup-product, is quadratic for every open subgroup U of G . Note that if $\text{im}(\theta) \subseteq 1 + p\mathbb{Z}_p$ and $p \neq 2$ then G acts trivially on $\mathbb{Z}_p(k)/p$. By the Rost-Voevodsky Theorem $(G_{\mathbb{K}}, \theta_{\mathbb{K}, p})$ is, indeed, Bloch-Kato.

For a profinite group G , let $O_p(G)$ denote the maximal closed normal pro- p subgroup of G . A p -oriented profinite group (G, θ) has two particular closed normal subgroups: the kernel $\ker(\theta)$ of θ , and the θ -center of (G, θ) , given by

$$Z_\theta(G) = \left\{ x \in O_p(\ker(\theta)) \mid gxg^{-1} = x^{\theta(g)} \text{ for all } g \in G \right\}. \quad (1.5)$$

As $Z_\theta(G)$ is contained in the center $Z(\ker(\theta))$ of $\ker(\theta)$, it is abelian. The p -oriented profinite group (G, θ) will be said to be θ -abelian, if $\ker(\theta) = Z_\theta(G)$ and if $Z_\theta(G)$ is torsion free. In particular, for such a p -oriented profinite group (G, θ) , G is a virtual pro- p group (i.e., G contains an open subgroup which is a pro- p group). Moreover, a θ -abelian pro- p group (G, θ) will be said to be *split* if $G \simeq Z_\theta(G) \rtimes \text{im}(\theta)$.

As $Z_\theta(G)$ is contained in $\ker(\theta)$, by definition, the canonical quotient $\bar{G} = G/Z_\theta(G)$ carries naturally a p -orientation $\theta: \bar{G} \rightarrow \mathbb{Z}_p^\times$, and one has the following short exact sequence of p -oriented profinite groups.

$$\{1\} \longrightarrow Z_\theta(G) \longrightarrow G \xrightarrow{\pi} \bar{G} \longrightarrow \{1\} \quad (1.6)$$

The following result can be seen as an analogue of the equal characteristic transition theorem (cf. [31, §II.4, Exercise 1(b), p. 86]) for cyclotomically p -oriented Bloch-Kato profinite groups.

THEOREM 1.2. *Let (G, θ) be a cyclotomically p -oriented Bloch-Kato profinite group. Then (1.6) splits, provided that $\text{cd}_p(G) < \infty$, and one of the following conditions hold:*

- (i) G is a pro- p group,
- (ii) (G, θ) is an oriented virtual pro- p group (see §4),
- (iii) $(\bar{G}, \bar{\theta})$ is cyclotomically p -oriented and Bloch-Kato.

In the case that (G, θ) is the maximal pro- p Galois group of a field \mathbb{K} containing a primitive p^{th} -root of unity endowed with the p -orientation induced by $\theta_{\mathbb{K}, p}$, $Z_\theta(G)$ is the inertia group of the maximal p -henselian valuation of \mathbb{K} (cf. Remark 7.8).

Note that the 2-oriented pro-2 group $(C_2 \times \mathbb{Z}_2, \theta)$ may be θ -abelian, but θ is never 1-cyclotomic (cf. Proposition 6.5). As a consequence, in a cyclotomically 2-oriented pro-2 group every element of order 2 is self-centralizing.

For p odd it was shown in [25] that a Bloch-Kato pro- p group G satisfies a strong form of *Tits alternative*, i.e., either G contains a closed non-abelian free pro- p subgroup, or there exists a p -orientation $\theta: G \rightarrow \mathbb{Z}_p^\times$ such that G is θ -abelian. In Subsection 7.1 we extend this result to pro-2 groups with a cyclotomic orientation, i.e., one has the following analogue of R. Ware's theorem (cf. [38]) for cyclotomically oriented Bloch-Kato pro- p groups (cf. Fact 7.4).

THEOREM 1.3. *Let (G, θ) be a cyclotomically p -oriented Bloch-Kato pro- p group. If $p = 2$ assume further that $\text{im}(\theta) \subseteq 1 + 4\mathbb{Z}_2$. Then one — and only one — of the following cases hold:*

- (i) G contains a closed non-abelian free pro- p subgroup; or
- (ii) G is θ -abelian.

It should be mentioned that for $p = 2$ the additional hypothesis is indeed necessary (cf. Remark 5.8). The class of cyclotomically p -oriented Bloch-Kato profinite groups is closed with respect to several constructions.

THEOREM 1.4. (a) *The inverse limit of an inverse system of cyclotomically p -oriented Bloch-Kato profinite groups with surjective structure maps is a cyclotomically p -oriented Bloch-Kato profinite group (cf. Corollary 3.3 and Corollary 3.6).*

- (b) *The free profinite (resp. pro- p) product of two cyclotomically p -oriented Bloch-Kato profinite (resp. pro- p) groups is a cyclotomically p -oriented Bloch-Kato profinite (resp. pro- p) group (cf. Theorem 3.14).*
- (c) *The fibre product of a cyclotomically p -oriented Bloch-Kato profinite group (G_1, θ_1) with a split θ_2 -abelian profinite group (G_2, θ_2) is a cyclotomically p -oriented Bloch-Kato profinite group (cf. Theorem 3.11 and Theorem 3.13).*
- (d) *The quotient of a cyclotomically p -oriented Bloch-Kato profinite group (G, θ) with respect to a closed normal subgroup $N \subseteq G$ satisfying $N \subseteq \ker(\theta)$ and N a p -perfect group is a cyclotomically p -oriented Bloch-Kato profinite group (cf. Proposition 4.6).*

Some time ago I. Efrat (cf. [7–9]) has formulated the so-called *elementary type conjecture* concerning the structure of finitely generated pro- p groups occurring as maximal pro- p quotients of an absolute Galois group. His conjecture can be

reformulated in the class of cyclotomically p -oriented Bloch-Kato pro- p groups. Such a p -oriented pro- p group (G, θ) is said to be *indecomposable* if $Z_\theta(G) = \{1\}$ and if G is not a proper free pro- p product. A positive answer to the following question would settle the elementary type conjecture affirmatively.

QUESTION 1.5. *Let (G, θ) be a finitely generated, torsion free, indecomposable, cyclotomically oriented Bloch-Kato pro- p group. Does this imply that G is a Poincaré duality pro- p group of dimension $\text{cd}_p(G) \leq 2$?*

The paper is organized as follows. In § 2 we give some equivalent definitions for cyclotomic p -orientations. In § 3 we study some operations of profinite groups (inverse limits, free products and fibre products) in relation with the properties of cyclotomicity and Bloch-Kato-ness, and we prove Theorem 1.4(a)-(b)-(c). In § 4 we study the quotients of cyclotomically p -oriented profinite groups over closed normal p -perfect subgroups — in particular, we introduce *oriented virtual pro- p groups* and we prove Theorem 1.4(d). In § 5 we study p -oriented profinite Poincaré duality groups. In § 6 we focus on the presence of torsion in cyclotomically 2-oriented pro-2 groups, and we prove that in a 1-cyclotomically 2-oriented pro-2 group every element of order 2 is self-centralizing (see Proposition 6.5). In § 7 we focus on the structure of cyclotomically p -oriented Bloch-Kato pro- p groups: we prove Theorems 1.2 and 1.3, and show that in many cases the θ -center is the maximal abelian closed normal subgroup (cf. Theorem 7.7).

2 ABSOLUTE GALOIS GROUPS AND CYCLOTOMIC p -ORIENTATIONS

Throughout the paper, we study profinite groups with a cyclotomic module $\mathbb{Z}_p(1)$. In contrast to [5, § 14], we refer to the associated representation $\theta: G \rightarrow \mathbb{Z}_p^\times$, rather than to the module itself. As we study several subgroups of G associated to this cyclotomic module $\mathbb{Z}_p(1)$, like $\ker(\theta)$ and $Z_\theta(G)$, this choice of notation turns out to be convenient for our purposes. We follow the convention as established in [25, 26] and call such representations “ p -orientations”.¹ In the case that G is a pro- p group, the couple (G, θ) was called a *cyclotomic pro- p pair*, in [9, § 3].

2.1 THE CONNECTING HOMOMORPHISM δ^k

Let G be a profinite group, and let $\theta: G \rightarrow \mathbb{Z}_p^\times$ be a p -orientation of G . For every $k \geq 0$ one has the short exact sequence of left $\mathbb{Z}_p[[G]]$ -modules

$$0 \longrightarrow \mathbb{Z}_p(k) \xrightarrow{p} \mathbb{Z}_p(k) \longrightarrow \mathbb{F}_p(k) \longrightarrow 0, \quad (2.1)$$

¹ For a Poincaré duality group G the representation associated to the *dualizing module* — which coincides with the cyclotomic module in the case of a Poincaré duality pro- p group of dimension 2 (cf. Theorem 5.7) — is sometimes also called the “orientation” of G .

which induces the long exact sequence in cohomology

$$\begin{array}{ccccccc}
 \cdots & \xrightarrow{\delta^{k-1}} & H_{\text{cts}}^k(G, \mathbb{Z}_p(k)) & \xrightarrow{p \cdot} & H_{\text{cts}}^k(G, \mathbb{Z}_p(k)) & \xrightarrow{\pi^k} & H_{\text{cts}}^k(G, \mathbb{F}_p(k)) \\
 & & & & & \searrow \delta^k & \\
 & & & & & & H_{\text{cts}}^{k+1}(G, \mathbb{Z}_p(k)) & \xrightarrow{p \cdot} & H_{\text{cts}}^{k+1}(G, \mathbb{Z}_p(k)) & \xrightarrow{\pi^{k+1}} & \cdots
 \end{array} \tag{2.2}$$

with connecting homomorphism δ^k (cf. [34, §2]). In particular, δ^k is trivial if, and only if, multiplication by p on $H_{\text{cts}}^{k+1}(G, \mathbb{Z}_p(k))$ is a monomorphism. This is equivalent to $H_{\text{cts}}^{k+1}(G, \mathbb{Z}_p(k))$ being torsion free. Therefore, one concludes the following:

PROPOSITION 2.1. *Let (G, θ) be a p -oriented profinite group. For $k \geq 1$ and $U \subseteq G$ an open subgroup the following are equivalent.*

- (i) *The map (1.2) is surjective for every $n \geq 1$.*
- (ii) *The map $\pi^k: H_{\text{cts}}^k(U, \mathbb{Z}_p(k)) \rightarrow H^k(U, \mathbb{F}_p(k))$ is surjective.*
- (iii) *The connecting homomorphism $\delta^k: H^k(U, \mathbb{F}_p(k)) \rightarrow H_{\text{cts}}^{k+1}(U, \mathbb{Z}_p(k))$ is trivial.*
- (iv) *The \mathbb{Z}_p -module $H_{\text{cts}}^{k+1}(U, \mathbb{Z}_p(k))$ is torsion free.*

Proof. By the long exact sequence (2.2), the equivalences between (ii), (iii) and (iv) are straightforward. For $m \geq n \geq 1$ let $\pi_{m,n}^k$ denote the natural maps

$$\pi_{m,n}^k: H^k(U, \mathbb{Z}_p(k)/p^m) \longrightarrow H^k(U, \mathbb{Z}_p(k)/p^n)$$

(if $m = \infty$ we set $p^\infty = 0$). If condition (i) holds then the system $(H^k(U, \mathbb{Z}_p/p^n), \pi_{m,n}^k)$ satisfies the Mittag-Leffler property. In particular,

$$H^k(U, \mathbb{Z}_p(k)) \simeq \varprojlim_{n \geq 1} H^k(U, \mathbb{Z}_p(k)/p^n)$$

(cf. [28] and [23, Thm. 2.7.5]). Thus $\pi^k = \pi_{n,1}^k \circ \pi_{\infty,n}^k$ is surjective if, and only if, $\pi_{n,1}^k$ is surjective for every $n \geq 1$. Conversely, if π^k is surjective then $\pi^k = \pi_{n,1}^k \circ \pi_{\infty,n}^k$ yields the surjectivity of $\pi_{n,1}^k$ for every n . □

2.2 PROFINITE GROUPS OF COHOMOLOGICAL p -DIMENSION AT MOST 1

Let G be a profinite group, and let $\theta: G \rightarrow \mathbb{Z}_p^\times$ be a p -orientation of G . Then

$$H_{\text{cts}}^1(G, \mathbb{Z}_p(0)) = \text{Hom}_{\text{grp}}(G, \mathbb{Z}_p) \tag{2.3}$$

is a torsion free abelian group for every profinite group G , i.e., θ is 0-cyclotomic. If G is of cohomological p -dimension less or equal to 1, then $H_{\text{cts}}^{m+1}(G, \mathbb{Z}_p(m)) =$

0 for all $m \geq 1$ showing that θ is cyclotomic. Moreover, $H^\bullet(G, \hat{\theta})$ is a quadratic \mathbb{F}_p -algebra for every profinite group with $\text{cd}_p(G) \leq 1$ and for any p -orientation $\theta: G \rightarrow \mathbb{Z}_p^\times$. If G is of cohomological p -dimension less or equal to 1, one has $\text{cd}_p(C) \leq 1$ for every closed subgroup C of G (cf. [31, §I.3.3, Proposition 14]). Thus one has the following.

FACT 2.2. *Let G be a profinite group with $\text{cd}_p(G) \leq 1$, and let $\theta: G \rightarrow \mathbb{Z}_p^\times$ be a p -orientation for G . Then (G, θ) is Bloch-Kato and θ is cyclotomic.*

2.3 THE m^{th} -NORM RESIDUE SYMBOL

Throughout this subsection we fix a field \mathbb{K} and a separable closure $\bar{\mathbb{K}}$ of \mathbb{K} . For $p \neq \text{char}(\mathbb{K})$, $\mu_{p^\infty}(\bar{\mathbb{K}})$ is a divisible abelian group. By construction, one has a canonical isomorphism

$$\varprojlim_{k \geq 0} (\mu_{p^\infty}(\bar{\mathbb{K}}), p^k) \simeq \mathbb{Z}_p(1) \otimes_{\mathbb{Z}} \mathbb{Q}_p = \mathbb{Q}_p(1) \quad (2.4)$$

and a short exact sequence $0 \rightarrow \mathbb{Z}_p(1) \rightarrow \mathbb{Q}_p(1) \rightarrow \mu_{p^\infty}(\bar{\mathbb{K}}) \rightarrow 0$ of topological left $\mathbb{Z}_p[[G_{\mathbb{K}}]]$ -modules, where $G_{\mathbb{K}} = \text{Gal}(\bar{\mathbb{K}}/\mathbb{K})$ is the absolute Galois group of \mathbb{K} .

Let $K_m^M(\mathbb{K})$, $m \geq 0$, denote the m^{th} -Milnor K -group of \mathbb{K} (cf. [10, §24.3]). For $p \neq \text{char}(\mathbb{K})$, J. Tate constructed in [34] a homomorphism of abelian groups

$$h_m(\mathbb{K}): K_m^M(\mathbb{K}) \longrightarrow H_{\text{cts}}^m(G_{\mathbb{K}}, \mathbb{Z}_p(m)), \quad (2.5)$$

the so-called m^{th} -norm residue symbol. Let $K_m^M(\mathbb{K})/p = K_m^M(\mathbb{K})/pK_m^M(\mathbb{K})$. Around ten years later S. Bloch and K. Kato conjectured in [1] that the induced map

$$h_m(\mathbb{K})/p: K_m^M(\mathbb{K})/p \longrightarrow H^m(G_{\mathbb{K}}, \mathbb{F}_p(m)) \quad (2.6)$$

is an isomorphism for all fields \mathbb{K} , $\text{char}(\mathbb{K}) \neq p$, and for all $m \geq 0$. This conjecture has been proved by V. Voevodsky and M. Rost with a “patch” of C. Weibel (cf. [29, 36, 40]). In particular, since $K_\bullet^M(\mathbb{K})/p$ is a quadratic \mathbb{F}_p -algebra and as $h_\bullet(\mathbb{K})/p$ is a homomorphism of algebras, this implies that the absolute Galois group of a field \mathbb{K} is Bloch-Kato (cf. [10, §23.4]). The Rost-Voevodsky Theorem has also the following consequence.

PROPOSITION 2.3. *Let \mathbb{K} be a field, let $G_{\mathbb{K}}$ denote its absolute Galois group, and let $\theta_{\mathbb{K}, p}: G_{\mathbb{K}} \rightarrow \mathbb{Z}_p^\times$ denote its canonical p -orientation. Then $\theta_{\mathbb{K}, p}$ is cyclotomic.*

Although Proposition 2.3 might be well known to specialists, we add a short proof of it. By Proposition 2.1, Proposition 2.3 in combination with Theorem 1.4-(d) is equivalent to [5, Prop. 14.19].

Proof of Proposition 2.3. If $\text{char}(\mathbb{K}) = p$, then $\text{cd}_p(G_{\mathbb{K}}) \leq 1$ (cf. [31, §II.2.2, Proposition 3]), and the p -orientation $\theta_{\mathbb{K}, p}$ is cyclotomic by Fact 2.2. So we

may assume that $\text{char}(\mathbb{K}) \neq p$. In the commutative diagram

$$\begin{array}{ccccccc}
 K_k^M(\mathbb{K}) & \xrightarrow{p} & K_k^M(\mathbb{K}) & \xrightarrow{\pi} & K_k^M(\mathbb{K})/p & \longrightarrow & 0 \\
 \downarrow h_k & & \downarrow h_k & & \downarrow (h_k)/p & & \\
 H_{\text{cts}}^k(G_{\mathbb{K}}, \mathbb{Z}_p(k)) & \xrightarrow{p} & H_{\text{cts}}^k(G_{\mathbb{K}}, \mathbb{Z}_p(k)) & \xrightarrow{\alpha} & H^k(G_{\mathbb{K}}, \mathbb{F}_p(k)) & \xrightarrow{\beta} & H_{\text{cts}}^{k+1}(G_{\mathbb{K}}, \mathbb{Z}_p(k))
 \end{array}
 \tag{2.7}$$

the map π is surjective, and $(h_k)/p$ is an isomorphism. Hence α must be surjective, and thus $\beta = 0$, i.e., $p: H_{\text{cts}}^{k+1}(G_{\mathbb{K}}, \mathbb{Z}_p(k)) \rightarrow H_{\text{cts}}^{k+1}(G_{\mathbb{K}}, \mathbb{Z}_p(k))$ is an injective homomorphism of \mathbb{Z}_p -modules. Thus $H_{\text{cts}}^{k+1}(G_{\mathbb{K}}, \mathbb{Z}_p(k))$ must be p -torsion free. Any open subgroup U of $G_{\mathbb{K}}$ is the absolute Galois group of \mathbb{K}^U . Hence $\theta_{\mathbb{K},p}$ is cyclotomic, and this yields the claim. \square

Remark 2.4. Let \mathbb{K} be a number field, let S be a set of places containing all infinite places of \mathbb{K} and all places lying above p , and let $G_{\mathbb{K}}^S$ be the Galois group of $\overline{\mathbb{K}}^S/\mathbb{K}$, where $\overline{\mathbb{K}}^S/\mathbb{K}$ is the maximal extension of $\overline{\mathbb{K}}/\mathbb{K}$ which is unramified outside S . Then $\theta_{\mathbb{K},p}: G_{\mathbb{K}} \rightarrow \mathbb{Z}_p^\times$ induces a p -orientation $\theta_{\mathbb{K},p}^S: G_{\mathbb{K}}^S \rightarrow \mathbb{Z}_p^\times$. However, it is well known (cf. [23, Prop. 8.3.11(ii)]) that,

$$H^1(G_{\mathbb{K}}^S, \mathbb{I}_p(1)) \simeq H^1(G_{\mathbb{K}}^S, \mathcal{O}_{\mathbb{K}}^S)_{(p)} \simeq \text{cl}(\mathcal{O}_{\mathbb{K}}^S)_{(p)}
 \tag{2.8}$$

(for the definition of $\mathbb{I}_p(1)$ see §3), where $\text{cl}(\mathcal{O}_{\mathbb{K}}^S)$ denotes the *ideal class group* of the Dedekind domain $\mathcal{O}_{\mathbb{K}}^S$, and $\text{---}_{(p)}$ denotes the p -primary component. Hence $(G_{\mathbb{K}}^S, \theta_{\mathbb{K},p}^S)$ is in general not cyclotomic (cf. Proposition 3.1).

3 COHOMOLOGY OF p -ORIENTED PROFINITE GROUPS

A homomorphism $\phi: (G_1, \theta_1) \rightarrow (G_2, \theta_2)$ of two p -oriented profinite groups (G_1, θ_1) and (G_2, θ_2) is a continuous group homomorphism $\phi: G_1 \rightarrow G_2$ satisfying $\theta_1 = \theta_2 \circ \phi$.

Let (G, θ) be a p -oriented profinite group. For $k \in \mathbb{Z}$, put $\mathbb{Q}_p(k) = \mathbb{Z}_p(k) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$, and also $\mathbb{I}_p(k) = \mathbb{Q}_p(k)/\mathbb{Z}_p(k)$, i.e., $\mathbb{I}_p(k)$ is a discrete left G -module and — as an abelian group — a divisible p -torsion module.

Let $\mathbb{I}_p = \mathbb{Q}_p/\mathbb{Z}_p$, and let $\text{---}^* = \text{Hom}_{\mathbb{Z}_p}(\text{---}, \mathbb{I}_p)$ denote the Pontryagin duality functor. Then $\mathbb{I}_p(k)^*$ is a profinite left $\mathbb{Z}_p[[G]]$ -module which is isomorphic to $\mathbb{Z}_p(-k)$.

3.1 CRITERIA FOR CYCLOTOMICITY

The following proposition relates the continuous co-chain cohomology groups, Galois cohomology and the Galois homology groups as defined by A. Brumer in [3].

PROPOSITION 3.1. *Let (G, θ) be a p -oriented profinite group, let k be an integer, and let m be a non-negative integer. Then the following are equivalent:*

- (i) $H_{\text{cts}}^{m+1}(G, \mathbb{Z}_p(k))$ is torsion free;
- (ii) $H^m(G, \mathbb{I}_p(k))$ is divisible;
- (iii) $H_m(G, \mathbb{Z}_p(-k))$ is torsion free.

Proof. The equivalence (i) \Leftrightarrow (ii) is a direct consequence of [34, Prop. 2.3], and (ii) \Leftrightarrow (iii) follows from [33, (3.4.5)]. \square

The direct limit of divisible p -torsion modules is a divisible p -torsion module. From this fact — and Proposition 3.1 — one concludes the following.

COROLLARY 3.2. *Let (G, θ) be a cyclotomically p -oriented profinite group. Then $H^m(C, \mathbb{I}_p(m))$ is divisible for all $m \geq 0$ and all C closed in G .*

Proof. It suffices to show (ii) \Rightarrow (i). Let C be a closed subgroup of G . Then $H^m(C, \mathbb{I}_p(m)) \simeq \varinjlim_{U \in \mathfrak{B}_C} H^m(U, \mathbb{I}_p(m))$, where \mathfrak{B}_C denotes the set of all open subgroups of G containing C (cf. [31, §I.2.2, Proposition 8]). Hence Proposition 3.1 yields the claim. \square

In combination with [3, Corollary 4.3(ii)], Proposition 3.1 implies the following.

COROLLARY 3.3. *Let (I, \preceq) be a directed set, let (G, θ) be a p -oriented profinite group, and let $(N_i)_{i \in I}$ be a family of closed normal subgroups of G satisfying $N_j \subseteq N_i \subseteq \ker(\theta)$ for $i \preceq j$ such that $\bigcap_{i \in I} N_i = \{1\}$ and the induced p -orientation $\theta_i: G/N_i \rightarrow \mathbb{Z}_p^\times$ is cyclotomic for all $i \in I$. Then $\theta: G \rightarrow \mathbb{Z}_p^\times$ is cyclotomic.*

Proof. Let $U \subseteq G$ be an open subgroup of G . Hypothesis (iii) implies that the group $H_m(UN_i/N_i, \mathbb{Z}_p(-m))$ is torsion free for all $i \in I$ (cf. Proposition 3.1). Thus, by [3, Corollary 4.3(ii)], $H_m(U, \mathbb{Z}_p(-m))$ is torsion free, and hence, by Proposition 3.1, $\theta: G \rightarrow \mathbb{Z}_p^\times$ is a cyclotomic p -orientation. \square

3.2 THE MOD- p COHOMOLOGY RING

An \mathbb{N}_0 -graded \mathbb{F}_p -algebra $\mathbf{A} = \coprod_{k \geq 0} \mathbf{A}_k$ is said to be *anti-commutative* if for $x \in \mathbf{A}_s$ and $y \in \mathbf{A}_t$ one has $y \cdot x = (-1)^{st} \cdot x \cdot y$. E.g., if V is an \mathbb{F}_p -vector space, the *exterior algebra* $\Lambda_\bullet(V)$ (cf. [18, Chapter 4]) is an \mathbb{N}_0 -graded anti-commutative \mathbb{F}_p -algebra. Moreover, if G is a profinite group, then its cohomology ring $H^\bullet(G, \mathbb{F}_p)$ is an \mathbb{N}_0 -graded anti-commutative \mathbb{F}_p -algebra (cf. [23, Prop. 1.4.4]).

Let $\mathbf{T}(V) = \coprod_{k \geq 0} V^{\otimes k}$ denote the *tensor algebra* generated by the \mathbb{F}_p -vector space V . A \mathbb{N}_0 -graded associative \mathbb{F}_p -algebra \mathbf{A} is said to be *quadratic* if the canonical homomorphism $\eta^{\mathbf{A}}: \mathbf{T}(\mathbf{A}_1) \rightarrow \mathbf{A}$ is surjective, and

$$\ker(\eta^{\mathbf{A}}) = \mathbf{T}(\mathbf{A}_1) \otimes \ker(\eta_2^{\mathbf{A}}) \otimes \mathbf{T}(\mathbf{A}_1) \quad (3.1)$$

(cf. [24, § 1.2]). E.g., $\mathbf{A} = \Lambda_\bullet(V)$ is quadratic.

If \mathbf{A} and \mathbf{B} are anti-commutative \mathbb{N}_0 -graded \mathbb{F}_p -algebras, then $\mathbf{A} \otimes \mathbf{B}$ is again an anti-commutative \mathbb{N}_0 -graded \mathbb{F}_p -algebra, where

$$(x_1 \otimes y_1) \cdot (x_2 \otimes y_2) = (-1)^{s_2 t_1} \cdot (x_1 \cdot x_2) \otimes (y_1 \cdot y_2), \tag{3.2}$$

for $x_1 \in \mathbf{A}_{s_1}, x_2 \in \mathbf{A}_{s_2}, y_1 \in \mathbf{B}_{t_1}, y_2 \in \mathbf{B}_{t_2}$. In particular, if \mathbf{A} and \mathbf{B} are quadratic, then $\mathbf{A} \otimes \mathbf{B}$ is quadratic as well.

A direct set (I, \preceq) maybe considered as a small category with objects given by the set I and precisely one morphism $\iota_{i,j}$ for all $i \preceq j, i, j \in I$, i.e., $\iota_{i,i} = \text{id}_i$. One has the following.

FACT 3.4. *Let \mathbb{F} be a field, let (I, \preceq) be a direct system, and let $\mathbf{A}: (I, \preceq) \rightarrow \mathbb{F}\mathbf{qalg}$ be a covariant functor with values in the category of quadratic \mathbb{F} -algebras. Then $\mathbf{B} = \varinjlim_{i \in \mathbf{A}} \mathbf{A}(i)$ is a quadratic \mathbb{F} -algebra.*

Let (G, θ) be a p -oriented profinite group, and let $\widehat{\theta}: G \rightarrow \mathbb{F}_p^\times$ be the map induced by θ . If $\widehat{\theta} = \mathbf{1}_G$, then the mod- p cohomology ring of $H^\bullet(G, \widehat{\theta})$ coincides with $H^\bullet(G, \mathbb{F}_p)$ (see (1.4)), and hence it is anti-commutative. Furthermore, if $\widehat{\theta} \neq \mathbf{1}_G$ and $G^\circ = \ker(\widehat{\theta})$, restriction

$$\text{res}_{G, G^\circ}^\bullet: H^\bullet(G, \widehat{\theta}) \longrightarrow H^\bullet(G^\circ, \mathbb{F}_p) \tag{3.3}$$

is an injective homomorphism of \mathbb{N}_0 -graded algebras. Hence the mod- p cohomology ring $H^\bullet(G, \theta)$ is anti-commutative. In particular, if $M_{(k)}$ denotes the homogeneous component of the left $\mathbb{F}_p[G/G^\circ]$ -module M , on which G/G° acts by $\widehat{\theta}^k$, the Hochschild-Serre spectral sequence (cf. [23, § II.4, Exercise 4(ii)]) shows that

$$H^k(G, \widehat{\theta}) = H^k(G^\circ, \mathbb{F}_p)_{(-k)}. \tag{3.4}$$

From [31, §I.2.2, Prop. 8] and Fact 3.4 one concludes the following.

COROLLARY 3.5. *Let (G, θ) be a p -oriented profinite group which is Bloch-Kato. Then $H^\bullet(C, \widehat{\theta}|_C)$ is quadratic for all C closed in G .*

COROLLARY 3.6. *Let (I, \preceq) be a directed set, let (G, θ) be a p -oriented profinite group, and let $(N_i)_{i \in I}$ be a family of closed normal subgroups of G , $N_j \subseteq N_i \subseteq \ker(\theta)$ for $i \preceq j$, such that $\bigcap_{i \in I} N_i = \{1\}$ and $(G/N_i, \widehat{\theta}_{N_i})$ is Bloch-Kato. Then (G, θ) is Bloch-Kato.*

Remark 3.7. Let G be a pro- p group with minimal presentation

$$G = \langle x_1, \dots, x_d \mid [x_1, x_2][[x_3, x_4], x_5] = 1 \rangle,$$

with $d \geq 5$. In [22, Ex. 7.3] and [21, § 4.3] it is shown that G does not occur as maximal pro- p Galois group of a field containing a primitive p^{th} -root of unity, relying on the properties of Massey products. It would be interesting to know whether G admits a cyclotomic p -orientation $\theta: G \rightarrow \mathbb{Z}_p^\times$ such that (G, θ) is Bloch-Kato. By Theorem 1.1, a negative answer would provide a ‘‘Massey-free’’ proof of the aforementioned fact.

3.3 FIBRE PRODUCTS

Let $(G_1, \theta_1), (G_2, \theta_2)$ be p -oriented profinite groups. The fibre product $(G, \theta) = (G_1, \theta_1) \boxtimes (G_2, \theta_2)$ denotes the pull-back of the diagram

$$\begin{array}{ccc}
 G_1 & \xrightarrow{\theta_1} & \mathbb{Z}_p^\times \\
 \uparrow & \nearrow \theta & \uparrow \theta_2 \\
 G & \dashrightarrow & G_2
 \end{array} \tag{3.5}$$

Remark 3.8. By restricting to the suitable subgroups if necessary, for the analysis of a fibre product $(G, \theta) = (G_1, \theta_1) \boxtimes (G_2, \theta_2)$ one may assume that $\text{im}(\theta_1) = \text{im}(\theta_2)$. In particular, if (G_2, θ_2) is split θ_2 -abelian and $G_2 \simeq A \rtimes \text{im}(\theta_2)$ for some free abelian pro- p group A , then $G \simeq A \rtimes G_1$ with $gag^{-1} = a^{\theta_1(g)}$ for all $a \in A$ and $g \in G_1$.

FACT 3.9. *Let (G, θ) be a p -oriented profinite group, and let N be a finitely generated non-trivial torsion free closed subgroup of $Z_\theta(G)$, i.e., $N \simeq \mathbb{Z}_p(1)^r$ as left $\mathbb{Z}_p[[G]]$ -modules for some $r \geq 1$. Then for $k \geq 0$ one has*

$$H^1(N, \mathbb{I}_p(k)) \simeq \mathbb{I}_p(k-1)^r \tag{3.6}$$

as left $\mathbb{Z}_p[[G]]$ -module.

The following property will be useful for the analysis of fibre products.

LEMMA 3.10. *Let $(G_1, \theta), (G_2, \theta_2)$ be cyclotomically p -oriented profinite groups, with (G_2, θ_2) split θ_2 -abelian and $Z = Z_{\theta_2}(G_2)$, and set*

$$(G, \theta) = (G_1, \theta_1) \boxtimes (G_2, \theta_2).$$

Let $\pi: G \rightarrow G_1$ be the canonical projection, and let $U \subseteq G$ be an open subgroup. Then $U \simeq (Z \cap U) \rtimes \pi(U)$.

Proof. Without loss of generality we may assume that $Z \simeq \mathbb{Z}_p$, so that $Z \cap U = Z^{p^k}$ for some $k \geq 0$. It suffices to show that there exists an open subgroup U_1 of U satisfying $Z \cap U_1 = \{1\}$ and $\pi(U_1) = \pi(U)$.

By choosing a section $\sigma: G_1 \rightarrow G$ (see Remark 3.8), one has a continuous homomorphism $\tau = \sigma \circ \pi: G \rightarrow G_1$ and a continuous function $\eta: G \rightarrow Z$ such that each $g \in G$ can be uniquely written as $g = \eta(g) \cdot \tau(g)$. In particular, for $h, h_1, h_2 \in U$ and $z \in Z \cap U = Z^{p^k}$ one has

$$\eta(z \cdot h) = z \cdot \eta(h) \quad \text{and} \quad \eta(h_1 \cdot h_2) = \eta(h_1) \cdot {}^{h_1}\eta(h_2). \tag{3.7}$$

Let $\eta_U = \chi \circ \eta|_U$, where $\chi: Z \rightarrow Z/Z^{p^k}$ is the canonical projection. By (3.7), η_U defines a crossed-homomorphism $\tilde{\eta}_U: \bar{U} \rightarrow Z/Z^{p^k}$, where $\bar{U} = U/Z^{p^k}$. As \bar{U} is canonically isomorphic to an open subgroup of G_1 , $(\bar{U}, \theta_1|_{\bar{U}})$ is cyclotomically

p -oriented. (Note that $Z \simeq \mathbb{Z}_p(1)$ as $\mathbb{Z}_p[[U]]$ -modules.) Hence, $H^1_{\text{cts}}(\bar{U}, \mathbb{Z}_p(1)) \rightarrow H^1(\bar{U}, \mathbb{Z}_p(1)/p^k)$ is surjective by Proposition 2.1, and the snake lemma applied to the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{B}^1(\bar{U}, Z) & \longrightarrow & \mathcal{Z}^1(\bar{U}, Z) & \longrightarrow & H^1(\bar{U}, \mathbb{Z}_p(1)) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{B}^1(\bar{U}, Z/Z^{p^k}) & \longrightarrow & \mathcal{Z}^1(\bar{U}, Z/Z^{p^k}) & \longrightarrow & H^1(\bar{U}, \mathbb{Z}_p(1)/p^k) \longrightarrow 0 \end{array} \quad (3.8)$$

where the left-side and right-side vertical arrows are surjective, shows that $\mathcal{Z}^1(\bar{U}, Z) \rightarrow \mathcal{Z}^1(\bar{U}, Z/Z^{p^k})$ is surjective. Thus there exists $\eta_\circ \in \mathcal{Z}^1(\bar{U}, Z)$ such that $\tilde{\eta}_U = \chi \circ \eta_\circ$. It is straightforward to verify that $U_1 = \{\eta_\circ(\bar{h}) \cdot \sigma(\bar{h}) \mid \bar{h} \in \bar{U}\}$ is an open subgroup of G_1 satisfying the requirements. \square

THEOREM 3.11. *Let (G_1, θ_1) be a cyclotomically p -oriented profinite group, and let (G_2, θ_2) be split θ_2 -abelian. Then $(G_1, \theta_1) \boxtimes (G_2, \theta_2)$ is cyclotomically p -oriented.*

Remark 3.12. (a) If p is odd, then every θ -abelian profinite group (G, θ) is split. However, a 2-oriented θ -abelian profinite group (G, θ) is split if, and only if, it is cyclotomically 2-oriented (cf. Proposition 6.7).

(b) If (G, θ) is θ -abelian and $H \subseteq G$ is a closed subgroup, then $(H, \theta|_H)$ is also θ -abelian.

Proof of Theorem 3.11. Put $(G, \theta) = (G_1, \theta_1) \boxtimes (G_2, \theta_2)$ and $Z = Z_{\theta_2}(G_2)$. We may also assume that $\text{im}(\theta_1) = \text{im}(\theta_2)$. As (G_2, θ_2) is split θ_2 -abelian, one has $G = Z \rtimes G_1$.

We first show the claim for $Z \simeq \mathbb{Z}_p$. Let U be an open subgroup of G . By Lemma 3.10, $(U, \theta|_U) \simeq (U_1, \theta_1) \boxtimes (U_2, \theta_2)$ where U_1 is isomorphic to an open subgroup of G_1 and (U_2, θ_2) is split θ_2 -abelian with $N = \ker(\theta_2)$ open in Z . As $\text{cd}_p(N) = 1$, one has $H^m(N, \mathbb{I}_p(k)) = 0$ for $m \geq 2$ and $k \geq 0$. Therefore, the E_2 -term of the Hochschild-Serre spectral sequence associated to the short exact sequence of profinite groups

$$\{1\} \longrightarrow N \longrightarrow U \longrightarrow U_1 \longrightarrow \{1\} \quad (3.9)$$

and evaluated on the discrete $\mathbb{Z}_p[[U]]$ -module $\mathbb{I}_p(k)$, is concentrated on the first and the second row. In particular, $d_r^{s,t} = 0$ for $r \geq 3$. As (3.9) splits, and as $\mathbb{I}_p(k)$ is inflated from U_1 , one has $E_2^{s,0}(\mathbb{I}_p(k)) = E_\infty^{s,0}(\mathbb{I}_p(k))$ for $s \geq 0$ (cf. [23, Prop. 2.4.5]). Hence $d_2^{s,t} = 0$ for all $s, t \geq 0$, i.e., $E_2^{s,t}(\mathbb{I}_p(k)) = E_\infty^{s,t}(\mathbb{I}_p(k))$, and the spectral sequence collapses. Thus, using the isomorphism (3.6), for every $k \geq 1$ one has a short exact sequence

$$0 \longrightarrow H^k(U_1, \mathbb{I}_p(k)) \xrightarrow{\text{inf}} H^k(U, \mathbb{I}_p(k)) \longrightarrow H^{k-1}(U_1, \mathbb{I}_p(k-1)) \longrightarrow 0, \quad (3.10)$$

where the right- and left-hand side are divisible p -torsion modules. As such \mathbb{Z}_p -modules are injective, (3.10) splits showing that $H^k(U, \mathbb{I}_p(k))$ is p -divisible. Therefore, by Proposition 3.1, (G, θ) is cyclotomic.

Thus, by induction the claim holds for all split θ_2 -abelian groups (G_2, θ_2) satisfying $\text{rk}(Z_{\theta_2}(G_2)) < \infty$. In general, as Z is a torsion free abelian pro- p group, there exists an inverse system $(Z_i)_{i \in I}$ of closed subgroups of Z such that Z/Z_i is torsion free, of finite rank, and $Z = \varprojlim_{i \in I} Z/Z_i$. Since Z_i is normal in G and

$$(G/Z_i, \bar{\theta}) \simeq (G_1, \theta_1) \boxtimes (G_2/Z_i, \bar{\theta}_2)$$

is cyclotomically p -oriented, Corollary 3.3 yields the claim. □

The following theorem can be seen as a generalization of a result of A. Wadsworth [37, Thm. 3.6].

THEOREM 3.13. *Let (G_i, θ_i) , $i = 1, 2$, be p -oriented profinite groups satisfying $\text{im}(\theta_1) = \text{im}(\theta_2)$. Assume further that (G_2, θ_2) is split θ_2 -abelian. Then for $(G, \theta) = (G_1, \theta_1) \boxtimes (G_2, \theta_2)$ one has that*

$$H^\bullet(G, \hat{\theta}) \simeq H^\bullet(G_1, \hat{\theta}_1) \otimes \Lambda_\bullet((\ker(\theta_2)/\ker(\theta_2)^p)^*). \tag{3.11}$$

Moreover, if (G_1, θ_1) is Bloch-Kato, then (G, θ) is Bloch-Kato.

Proof. Assume first that $d(Z_{\theta_2}(G_2))$ is finite. If $d(Z_{\theta_2}(G_2)) = 1$ then one obtains the isomorphism (3.11) from [37, Thm. 3.1], which uses the Hochschild-Serre spectral sequence associated to the short exact sequence of profinite groups

$$\{1\} \longrightarrow Z_{\theta_2}(G_2) \longrightarrow G \longrightarrow G/Z_{\theta_2}(G_2) \longrightarrow \{1\}$$

and evaluated on the discrete $\mathbb{Z}_p[[G]]$ -module $\mathbb{F}_p(k)$, to compute $H^\bullet(G, \hat{\theta})$. If $d(Z_{\theta_2}(G_2)) > 1$, then applying induction on $d(Z_{\theta_2}(G_2))$ yields the isomorphism (3.11). Finally, if $Z_{\theta_2}(G_2)$ is not finitely generated, then a limit argument similar to the one used in the proof Theorem 3.11 and Corollary 3.6 yield the claim. □

3.4 COPRODUCTS

For two profinite groups G_1 and G_2 let $G = G_1 \amalg G_2$ denote the coproduct (or free product) in the category of profinite groups (cf. [27, § 9.1]). In particular, if (G_1, θ_1) and (G_2, θ_2) are two p -oriented profinite groups, the p -orientations θ_1 and θ_2 induce a p -orientation $\theta: G \rightarrow \mathbb{Z}_p^\times$ via the universal property of the free product. Thus, we may interpret \amalg as the coproduct in the category of p -oriented profinite groups (cf. [9, §3]). The same applies to \amalg^p — the coproduct in the category of pro- p groups.

THEOREM 3.14. *Let (G_1, θ_1) and (G_2, θ_2) be two cyclotomically p -oriented profinite groups. Then their coproduct $(G, \theta) = (G_1, \theta_1) \amalg (G_2, \theta_2)$ is cyclotomically oriented. Moreover, if (G_1, θ_1) and (G_2, θ_2) are Bloch-Kato, then (G, θ) is Bloch-Kato.*

Proof. Let $(U, \theta|_U)$ be an open subgroup of (G, θ) . Then, by the Kurosh subgroup theorem (cf. [27, Thm. 9.1.9]),

$$U \simeq \prod_{s \in \mathcal{S}_1} ({}^s G_1 \cap U) \amalg \prod_{t \in \mathcal{S}_2} ({}^t G_2 \cap U) \amalg F, \tag{3.12}$$

where ${}^y G_i = yG_i y^{-1}$ for $y \in G$. The sets \mathcal{S}_1 and \mathcal{S}_2 are sets of representatives of the double cosets $U \backslash G / G_1$ and $U \backslash G / G_2$, respectively. In particular, the sets \mathcal{S}_1 and \mathcal{S}_2 are finite, and F is a free profinite subgroup of finite rank.

Put $U_s = {}^s G_1 \cap U$ for all $s \in \mathcal{S}_1$, and $V_t = {}^t G_2 \cap U$ for all $t \in \mathcal{S}_2$. By [23, Thm. 4.1.4], one has an isomorphism

$$H^k(U, \mathbb{I}_p(k)) \simeq \bigoplus_{s \in \mathcal{S}_1} H^k(U_s, \mathbb{I}_p(k)) \oplus \bigoplus_{t \in \mathcal{S}_2} H^k(V_t, \mathbb{I}_p(k)), \tag{3.13}$$

for $k \geq 2$, and an exact sequence

$$M \xrightarrow{\alpha} H^1(U, \mathbb{I}_p(1)) \longrightarrow M' \longrightarrow 0. \tag{3.14}$$

If (G_1, θ_1) and (G_2, θ_2) are cyclotomically p -oriented, then, by hypothesis and (3.13), $H^k(U, \mathbb{I}_p(k))$ is a divisible p -torsion module for $k \geq 2$. In (3.14), the module M is a homomorphic image of a p -divisible p -torsion module, and the module M' is the direct sum of p -divisible p -torsion modules, showing that $H^1(U, \mathbb{I}_p(1))$ is divisible. Hence, by Proposition 3.1 and Corollary 3.3, (G, θ) is cyclotomically p -oriented.

Assume that (G_1, θ_1) and (G_2, θ_2) are Bloch-Kato. Then — for U as in (3.12) — one has by (3.13) and (3.14) that

$$H^\bullet(U, \widehat{\theta}|_U) \simeq \mathbf{A} \oplus \bigoplus_{s \in \mathcal{S}_1} H^\bullet(U_s, \widehat{\theta}|_{U_s}) \oplus \bigoplus_{t \in \mathcal{S}_2} H^\bullet(V_t, \widehat{\theta}|_{V_t}) \oplus H^\bullet(F, \widehat{\theta}|_F) \tag{3.15}$$

where \mathbf{A} is a quadratic algebra, and \oplus denotes the *direct sum* in the category of quadratic algebras (cf. [24, p. 55]). In particular, $H^\bullet(U, \widehat{\theta}|_U)$ is quadratic. \square

For pro- p groups one has also the following.

THEOREM 3.15. *Let (G_1, θ_1) and (G_2, θ_2) be two cyclotomically oriented pro- p groups. Then their coproduct $(G, \theta) = (G_1, \theta_1) \amalg^p (G_2, \theta_2)$ is cyclotomically oriented. Moreover, if (G_1, θ_1) and (G_2, θ_2) are Bloch-Kato, then (G, θ) is Bloch-Kato.*

Proof. The Kurosh subgroup theorem is also valid in the category of pro- p groups with \amalg^p replacing \amalg (cf. [27, Thm. 9.1.9]), and (3.13) and (3.14) hold also in this context (cf. [23, Thm. 4.1.4]). Hence the proof for cyclotomicity can be transferred verbatim. The Bloch-Kato property was already shown in [25, Thm. 5.2]. \square

4 ORIENTED VIRTUAL PRO- p GROUPS

We say that a p -oriented profinite group (G, θ) is an *oriented virtual pro- p group* if $\ker(\theta)$ is a pro- p group. In particular, G is a virtual pro- p group. Since \mathbb{Z}_2^\times is a pro-2 group, every oriented virtual pro-2 group is in fact a pro-2 group. For $p \neq 2$ let $\hat{\theta}: G \rightarrow \mathbb{F}_p^\times$ be the homomorphism induced by θ , and put $G^\circ = \ker(\hat{\theta})$. Then $G/G^\circ \simeq \text{im}(\hat{\theta})$ is a finite cyclic group of order co-prime to p . The profinite version of the Schur-Zassenhaus theorem (cf. [14, Lemma 22.10.1]) implies that the short exact sequence of profinite groups

$$\{1\} \longrightarrow G^\circ \longrightarrow G \xrightarrow{\hat{\theta}} \text{im}(\hat{\theta}) \longrightarrow \{1\} \tag{4.1}$$

$\longleftarrow \sigma \longrightarrow$

splits. Indeed, if $C \subseteq G$ is a p' -Hall subgroup of G , then $\pi|_C: C \rightarrow \text{im}(\hat{\theta})$ is an isomorphism, and $\sigma = (\pi|_C)^{-1}$ is a canonical section for $\hat{\theta}$.

Note that $\mathbb{Z}_p^\times = \mathbb{F}_p^\times \times \Xi_p$, where $\Xi_p = O_p(\mathbb{Z}_p^\times)$ is the pro- p Sylow subgroup of \mathbb{Z}_p^\times , and where we denoted by \mathbb{F}_p^\times also the image of the Teichmüller section $\tau: \mathbb{F}_p^\times \rightarrow \mathbb{Z}_p^\times$. Hence a p -orientation $\theta: G \rightarrow \mathbb{Z}_p^\times$ on G defines a homomorphism $\hat{\theta}: G \rightarrow \mathbb{F}_p^\times$ and also a homomorphism $\theta^\vee: G \rightarrow \Xi_p$. On the contrary a pair of continuous homomorphisms $(\hat{\theta}, \theta^\vee)$, where $\hat{\theta}: G \rightarrow \mathbb{F}_p^\times$ and $\theta^\vee: G \rightarrow \Xi_p$, defines a p -orientation $\theta: G \rightarrow \mathbb{Z}_p^\times$ given by $\theta(g) = \hat{\theta}(g) \cdot \theta^\vee(g)$ for $g \in G$.

FACT 4.1. *Let $\hat{\theta}: G \rightarrow \mathbb{F}_p^\times$, $\sigma: \text{im}(\hat{\theta}) \rightarrow G$ be homomorphisms of groups satisfying (4.1). A homomorphism $\theta^\circ: G^\circ \rightarrow \Xi_p$ defines a p -orientation $\theta: G \rightarrow \mathbb{Z}_p^\times$, provided for all $c \in \text{im}(\hat{\theta})$ and for all $g \in G^\circ$ one has*

$$\theta^\circ(\sigma(c) \cdot g \cdot \sigma(c)^{-1}) = \theta^\circ(g) \tag{4.2}$$

Proof. By (4.1), one has $G = G^\circ \rtimes_\beta \bar{\Sigma}$, where $\bar{\Sigma} = \text{im}(\hat{\theta})$, $\beta: \bar{\Sigma} \rightarrow \text{Aut}(G^\circ)$ and $\beta(c)$ is left conjugation by $\sigma(c)$ for $c \in \bar{\Sigma}$. Thus, by (4.2), the map $\theta^\vee: G \rightarrow \Xi_p$ given by $\theta^\vee(g, c) = \theta^\circ(g)$ is a continuous homomorphism of groups, and (ι, θ^\vee) , where $\iota: \bar{\Sigma} \rightarrow \mathbb{F}_p^\times$ is the canonical inclusion, defines a p -orientation of G . \square

Let (G, θ) be an oriented virtual pro- p group satisfying (4.1). As $\theta: G \rightarrow \mathbb{Z}_p^\times$ is a homomorphism onto an abelian group one has

$$\theta(c \cdot g \cdot c^{-1}) = \theta(g) \tag{4.3}$$

for all $c \in C = \text{im}(\sigma)$ and $g \in G$. Thus, if $i_c \in \text{Aut}(G)$ denotes left conjugation by $c \in C$, one has

$$\theta = \theta \circ i_c \tag{4.4}$$

for all $c \in C$.

4.1 ORIENTED $\bar{\Sigma}$ -VIRTUAL PRO- p GROUPS

From now on let p be odd, and fix a subgroup $\bar{\Sigma}$ of \mathbb{F}_p^\times . An oriented virtual pro- p group (G, θ) is said to be an oriented $\bar{\Sigma}$ -virtual pro- p group, if $\text{im}(\hat{\theta}) = \bar{\Sigma}$. Hence, by the previous subsection, for such a group one has a split short exact sequence

$$\{1\} \longrightarrow G^\circ \longrightarrow G \xrightarrow{\hat{\theta}} \bar{\Sigma} \longrightarrow \{1\} . \tag{4.5}$$

$\underbrace{\hspace{1.5cm}}_{\sigma}$

By abuse of notation, we consider from now on (G, θ, σ) as an oriented $\bar{\Sigma}$ -virtual pro- p group. As the following fact shows there is also an *alternative form* of a $\bar{\Sigma}$ -virtual pro- p group.

FACT 4.2. *Let $\bar{\Sigma}$ be a subgroup of \mathbb{F}_p^\times . Let Q be a pro- p group, let $\theta^\circ: Q \rightarrow \Xi_p$ be a continuous homomorphism, and let $\gamma_Q: \bar{\Sigma} \rightarrow \text{Aut}_c(Q)$ be a homomorphism of groups, where $\text{Aut}_c(_)$ is the group of continuous automorphisms, satisfying*

$$\theta^\circ(\gamma_Q(c)(q)) = \theta^\circ(q), \tag{4.6}$$

for all $q \in Q$ and $c \in \bar{\Sigma}$, then $(Q \rtimes_{\gamma_Q} \bar{\Sigma}, \theta, \iota)$ is an oriented $\bar{\Sigma}$ -virtual pro- p group, where $\iota: \bar{\Sigma} \rightarrow Q \rtimes_{\gamma_Q} \bar{\Sigma}$ is the canonical map, and $\theta: Q \rtimes_{\gamma_Q} \bar{\Sigma} \rightarrow \mathbb{Z}_p^\times$ is the homomorphism induced by θ° (cf. Fact 4.1).

If $(G_1, \theta_1, \sigma_1)$ and $(G_2, \theta_2, \sigma_2)$ are oriented $\bar{\Sigma}$ -virtual pro- p groups, a continuous group homomorphism $\phi: G_1 \rightarrow G_2$ is said to be a morphism of $\bar{\Sigma}$ -virtual pro- p groups, if $\sigma_2 = \phi \circ \sigma_1$ and $\theta_1 = \theta_2 \circ \phi$. Similarly, if $(Q, \theta_Q^\circ, \gamma_Q)$ and $(R, \theta_R^\circ, \gamma_R)$ are $\bar{\Sigma}$ -virtual pro- p groups in alternative form (cf. Fact 4.2), the continuous group homomorphism $\phi: Q \rightarrow R$ is a homomorphisms of $\bar{\Sigma}$ -virtual pro- p groups provided $\theta_R \circ \phi = \theta_Q$ and if for all $c \in \bar{\Sigma}$ and for all $q \in Q$ one has that

$$\gamma_R(c)(\phi(q)) = \phi(\gamma_Q(c)(q)). \tag{4.7}$$

With this slightly more sophisticated set-up the category of $\bar{\Sigma}$ -virtual pro- p groups admits coproducts. In more detail, let $(Q, \theta_Q^\circ, \gamma_Q)$ and $(R, \theta_R^\circ, \gamma_R)$ be $\bar{\Sigma}$ -virtual pro- p groups in alternative form. Put $X = Q \amalg R$. Then for every element $c \in \bar{\Sigma}$ there exists an element $\delta(c) \in \text{Aut}(X)$ making the diagram

$$\begin{array}{ccccc} Q & \xrightarrow{\iota_1} & X & \xleftarrow{\iota_2} & R \\ \gamma_Q(c) \downarrow & & \downarrow \delta(c) & & \downarrow \gamma_R(c) \\ Q & \xrightarrow{\iota_1} & X & \xleftarrow{\iota_2} & R \end{array} \tag{4.8}$$

commute. Since Ξ_p is a pro- p group, there exists a continuous group homo-

morphism $\theta^\circ : X \rightarrow \Xi_p$ making the lower two rows of the diagram

$$\begin{array}{ccccc}
 Q & \xrightarrow{j_Q} & X & \xleftarrow{j_R} & R \\
 \downarrow \gamma_Q(c) & & \downarrow \delta(c) & & \downarrow \gamma_R(c) \\
 Q & \xrightarrow{j_Q} & X & \xleftarrow{j_R} & R \\
 \searrow \theta_Q^\circ & & \downarrow \theta^\circ & & \swarrow \theta_R^\circ \\
 & & \Xi_p & &
 \end{array} \tag{4.9}$$

commute. Since $\theta_{Q/R}^\circ = \theta_{Q/R}^\circ \circ \gamma_{Q/R}(c)$ for all $c \in \bar{\Sigma}$, one has $\theta^\circ = \theta^\circ \circ \delta(c)$ for all $c \in \bar{\Sigma}$. The commutativity of the diagram (4.9) yields that the group homomorphisms $j_Q : (Q, \theta_Q^\circ, \gamma_Q) \rightarrow (X, \theta^\circ, \delta)$ and $j_R : (R, \theta_R^\circ, \gamma_R) \rightarrow (X, \theta^\circ, \delta)$ are homomorphisms of oriented $\bar{\Sigma}$ -virtual pro- p groups in alternative form. Moreover, one has the following.

PROPOSITION 4.3. *The oriented $\bar{\Sigma}$ -virtual pro- p group $(X, \theta^\circ, \delta)$ together with the homomorphisms $j_Q : Q \rightarrow X$, and $j_R : R \rightarrow X$ is a coproduct in the category of oriented $\bar{\Sigma}$ -virtual pro- p groups.*

Proof. Let (H, θ_H, γ_H) be an oriented $\bar{\Sigma}$ -virtual pro- p group in alternative form, and let $\phi_Q : Q \rightarrow H$ and $\phi_R : R \rightarrow H$ be homomorphisms of oriented $\bar{\Sigma}$ -virtual pro- p groups in alternative form. Then there exists a unique homomorphism of pro- p groups $\phi : X \rightarrow H$ making the diagram concentrated on the second and third row of

$$\begin{array}{ccccc}
 Q & \xrightarrow{j_Q} & X & \xleftarrow{j_R} & R \\
 \downarrow \gamma_Q(c) & & \downarrow \delta(c) & & \downarrow \gamma_R(c) \\
 Q & \xrightarrow{j_Q} & X & \xleftarrow{j_R} & R \\
 \searrow \phi_Q & & \downarrow \phi & & \swarrow \phi_R \\
 & & H & & \\
 \searrow \theta_Q^\circ & & \downarrow \theta_H^\circ & & \swarrow \theta_R^\circ \\
 & & \Xi_p & &
 \end{array} \tag{4.10}$$

commute. Since $\phi_{Q/R} \circ \gamma_{Q/R}(c) = \gamma_H(c) \circ \phi_{Q/R}$ for all $c \in \bar{\Sigma}$, the uniqueness of ϕ implies that $\phi \circ \delta(c) = \gamma_H(c) \circ \phi$ for all $c \in \bar{\Sigma}$. As $\phi_Q : Q \rightarrow H$ and $\phi_R : R \rightarrow H$ are homomorphisms of $\bar{\Sigma}$ -virtual pro- p groups, one has that $\theta_{Q/R}^\circ = \theta_H^\circ \circ \phi_{Q/R}$. This implies that $(\theta_H^\circ \circ \phi) \circ j_{Q/R} = \theta_{Q/R}^\circ$, and from the construction of $\theta^\circ : X \rightarrow \Xi_p$ one concludes that $\theta^\circ = \theta_H^\circ \circ \phi$. This implies that ϕ is a homomorphism of oriented $\bar{\Sigma}$ -virtual pro- p groups. \square

Example 4.4. For $p = 3$ set $\bar{\Sigma} = \mathbb{F}_3^\times = \{1, s\}$. Then $(\mathbb{Z}_3^\times, \text{id}) \amalg^{\bar{\Sigma}} (\mathbb{Z}_3^\times, \text{id})$ is

isomorphic to $F \rtimes \bar{\Sigma}$, where $F = \langle x, y \rangle$ is a free pro-3 group of rank 2 and the induced isomorphism $s: F \rightarrow F$ satisfies $s(x) = x^{-1}$, $s(y) = y^{-1}$.

PROPOSITION 4.5. *Let (Q, θ_Q, γ_Q) be an oriented $\bar{\Sigma}$ -virtual pro- p group, and let Z be a normal $\bar{\Sigma}$ -invariant subgroup of Q isomorphic to \mathbb{Z}_p , which is not contained in the Frattini subgroup $\Phi(Q) = \text{cl}([Q, Q]Q^p)$ of Q . Then there exists a maximal closed subgroup M of Q which is $\bar{\Sigma}$ -invariant, such that $M \cdot Z = Q$ and $M \cap Z = \mathbb{Z}^p$.*

Proof. Let $\bar{Q} = Q/\Phi(Q)$. Then γ_Q induces a homomorphism $\bar{\gamma}_{\bar{Q}}: \bar{\Sigma} \rightarrow \text{Aut}_c(\bar{Q})$ making \bar{Q} a compact $\mathbb{F}_p[\bar{\Sigma}]$ -module. Let $\Omega = \text{Hom}_{\bar{\Sigma}}^c(\bar{Q}, \mathbb{F}_p)$, where \mathbb{F}_p denotes the finite field \mathbb{F}_p with canonical left $\bar{\Sigma}$ -action. By Pontryagin duality, one has $\bigcap_{\omega \in \Omega} \ker(\omega) = \{0\}$. Thus, by hypothesis, there exists $\psi \in \Omega$ such that $\psi|_Z \neq 0$. Hence $M = \ker(\psi)$ has the desired properties. \square

4.2 THE MAXIMAL ORIENTED VIRTUAL PRO- p QUOTIENT

For a prime p and a profinite group G we denote by $O^p(G)$ the closed subgroup of G generated by all Sylow pro- ℓ subgroups of G , $\ell \neq p$. In particular, $O^p(G)$ is p -perfect, i.e., $H^1(O^p(G), \mathbb{F}_p) = 0$, and one has the short exact sequence

$$\{1\} \longrightarrow O^p(G) \longrightarrow G \longrightarrow G(p) \longrightarrow \{1\},$$

where $G(p)$ denotes the maximal pro- p quotient of G . For a p -oriented profinite group (G, θ) , we denote by

$$G(\theta) = G/O^p(G^\circ)$$

the maximal p -oriented virtual pro- p quotient of G (for the definition of G° see the beginning of § 4). By construction, it carries naturally a p -orientation $\theta: G(\theta) \rightarrow \mathbb{Z}_p^\times$ inherited by G .

Note that if $\text{im}(\theta)$ is a pro- p group, then $G^\circ = G$, and $G(\theta) = G(p)$.

PROPOSITION 4.6. *Let (G, θ) be a p -oriented Bloch-Kato profinite group, and let $O \subseteq G$ be a p -perfect subgroup such that $O \subseteq \ker(\theta)$. Then the inflation map*

$$\text{inf}^k(M): H_{\text{cts}}^k(G/O, M) \longrightarrow H_{\text{cts}}^k(G, M), \tag{4.11}$$

is an isomorphism for all $k \geq 0$ and all $M \in \text{ob}(\mathbb{Z}_p[[G/O]]\mathbf{prf})$, where $\mathbb{Z}_p[[G/O]]\mathbf{prf}$ denotes the abelian category of profinite left $\mathbb{Z}_p[[G/O]]$ -modules.

Proof. As $O \subseteq \ker(\theta)$, $\mathbb{Z}_p(k)$ is a trivial $\mathbb{Z}_p[[O]]$ -module for every $k \in \mathbb{Z}$. Since O is p -perfect, and as the \mathbb{F}_p -algebra $H^\bullet(O, \mathbb{F}_p)$ is quadratic, $H^\bullet(O, \mathbb{F}_p)$ is 1-dimensional concentrated in degree 0. By Pontryagin duality, this is equivalent to $H_k(O, \mathbb{F}_p) = 0$ for all $k > 0$, where $H_k(O, _)$ denotes Galois homology as defined by A. Brumer in [3]. Thus, the long exact sequence in Galois homology implies that $H_k(O, \mathbb{Z}_p) = 0$ for all $k > 0$.

Let $(P_\bullet, \partial_\bullet, \varepsilon)$ be a projective resolution of the trivial left $\mathbb{Z}_p[[G]]$ -module in the category ${}_{\mathbb{Z}_p[[G]]}\mathbf{prf}$. For a projective left $\mathbb{Z}_p[[G]]$ -module $P \in \text{ob}({}_{\mathbb{Z}_p[[G]]}\mathbf{prf})$ define

$$\text{def}(P) = \text{def}_{G/O}^G(P) = \mathbb{Z}_p[[G/O]] \widehat{\otimes}_G P, \quad (4.12)$$

where $\widehat{\otimes}$ denotes the completed tensor product as defined in [3]. Then, by the Eckmann-Shapiro lemma in homology, one has that

$$H_k(\text{def}(P_\bullet), \text{def}(\partial_\bullet)) \simeq H_k(O, \mathbb{Z}_p). \quad (4.13)$$

Hence, by the previously mentioned remark, $(\text{def}(P_\bullet), \text{def}(\partial_\bullet))$ is a projective resolution of \mathbb{Z}_p in the category ${}_{\mathbb{Z}_p[[G/O]]}\mathbf{prf}$.

Let $M \in \text{ob}({}_{\mathbb{Z}_p[[G/O]]}\mathbf{prf})$. Then for every projective profinite left $\mathbb{Z}_p[[G]]$ -module P , one has a natural isomorphism

$$\text{Hom}_{G/O}(\text{def}(P), M) \simeq \text{Hom}_G(P, M). \quad (4.14)$$

Hence $\text{Hom}_{G/O}(\text{def}(P_\bullet), M)$ and $\text{Hom}_G(P_\bullet, M)$ are isomorphic co-chain complexes, and the induced maps in cohomology — which coincide with $\text{inf}^\bullet(M)$ — are isomorphisms. \square

COROLLARY 4.7. *Let (G, θ) be a p -oriented profinite group which is Bloch-Kato, respectively cyclotomically oriented. Then the maximal oriented virtual pro- p quotient $(G(\theta), \theta)$ is Bloch-Kato, respectively cyclotomically oriented.*

5 PROFINITE POINCARÉ DUALITY GROUPS AND p -ORIENTATIONS

5.1 PROFINITE POINCARÉ DUALITY GROUPS

Let G be a profinite group, and let p be a prime number. Then G is called a p -Poincaré duality group of dimension d , if

$$(PD_1) \quad \text{cd}_p(G) = d;$$

$$(PD_2) \quad |H_{\text{cts}}^k(G, A)| < \infty \text{ for every finite discrete left } G\text{-module } A \text{ of } p\text{-power order};$$

$$(PD_3) \quad H_{\text{cts}}^k(G, \mathbb{Z}_p[[G]]) = 0 \text{ for } k \neq d, \text{ and } H_{\text{cts}}^d(G, \mathbb{Z}_p[[G]]) \simeq \mathbb{Z}_p.$$

Although quite different at first glance, for a pro- p group our definition of p -Poincaré duality coincides with the definition given by J-P. Serre in [31, §I.4.5]. However, some authors prefer to omit the condition (PD_2) in the definition of a p -Poincaré duality group (cf. [23, Chap. III, §7, Definition 3.7.1]).

For a profinite p -Poincaré duality group G of dimension d the profinite right $\mathbb{Z}_p[[G]]$ -module $D_G = H_{\text{cts}}^d(G, \mathbb{Z}_p[[G]])$ is called the *dualizing module*. Since D_G is isomorphic to \mathbb{Z}_p as a pro- p group, there exists a unique p -orientation $\bar{\partial}_G: G \rightarrow \mathbb{Z}_p^\times$ such that for $g \in G$ and $z \in D_G$ one has

$$z \cdot g = z \cdot \bar{\partial}_G(g) = \bar{\partial}_G(g) \cdot z.$$

We call $\check{\delta}_G$ the *dualizing p -orientation*.

Let ${}^\times D_G$ denote the associated profinite left $\mathbb{Z}_p[[G]]$ -module, i.e., setwise ${}^\times D_G$ coincides with D_G and for $g \in G$ and $z \in {}^\times D_G$ one has

$$g \cdot z = z \cdot g^{-1} = \check{\delta}_G(g^{-1}) \cdot z.$$

For a profinite p -Poincaré duality group of dimension d the usual standard arguments (cf. [2, §VIII.10] for the discrete case) provide natural isomorphisms

$$\begin{aligned} \mathrm{Tor}_k^G(D_G, _)\simeq H_{\mathrm{cts}}^{d-k}(G, _), \\ \mathrm{Ext}_G^k({}^\times D_G, _)\simeq H_{d-k}(G, _), \end{aligned} \tag{5.1}$$

where $\mathrm{Tor}_\bullet^G(_, _)$ denotes the left derived functor of $_ \widehat{\otimes}_G _$, and $\mathrm{Ext}_G^\bullet(_, _)$ denotes the right derived functors of $\mathrm{Hom}_G(_, _)$ in the category $\mathbb{Z}_p[[G]]\mathbf{prf}$ (cf. [3]).

If A is a discrete left G -module which is also a p -torsion module, then A^* carries naturally the structure of a left (profinite) $\mathbb{Z}_p[[G]]$ -module (cf. [27, p. 171]). Then, by [31, § I.3.5, Proposition 17], Pontryagin duality and [33, (3.4.5)], one obtains for every finite discrete left $\mathbb{Z}_p[[G]]$ -module A of p -power order that

$$H_{\mathrm{cts}}^d(G, A) \simeq \mathrm{Hom}_G(A, I_G)^* \simeq \mathrm{Hom}_G(I_G^*, A^*)^* \simeq (I_G^*)^\times \widehat{\otimes}_G A, \tag{5.2}$$

where I_G denotes the discrete left dualizing module of G (cf. [31, §I.3.5]). In particular, by (5.1), $D_G \simeq (I_G^*)^\times$.

Example 5.1. Let $G_{\mathbb{K}}$ be the absolute Galois group of an ℓ -adic field \mathbb{K} . Then $G_{\mathbb{K}}$ satisfies p -Poincaré duality of dimension 2 for all prime numbers p . One has $I_G \simeq \mu_{p^\infty}(\overline{\mathbb{K}})$ (cf. [31, §II.5.2, Theorem 1]). Hence ${}^\times D_{G_{\mathbb{K}}} \simeq \mathbb{Z}_p(-1)$ with respect to the cyclotomic p -orientation $\theta_{\mathbb{K},p}: G_{\mathbb{K}} \rightarrow \mathbb{Z}_p^\times$, i.e., $\check{\delta}_{G_{\mathbb{K}}} = \theta_{\mathbb{K},p}$.

As we will see in the next proposition, the final conclusion in Example 5.1 is a consequence of a general property of Poincaré duality groups.

PROPOSITION 5.2. *Let G be a p -Poincaré duality group of dimension d , and let $\theta: G \rightarrow \mathbb{Z}_p^\times$ be a cyclotomic p -orientation of G . Then $\theta^{d-1} = \check{\delta}_G$ and ${}^\times D_G \simeq \mathbb{Z}_p(1-d)$.*

Proof. By (5.1) and the hypothesis, $H_{\mathrm{cts}}^d(G, \mathbb{Z}_p(d-1)) \simeq D_G \widehat{\otimes} \mathbb{Z}_p(d-1)$ is torsion free, and hence isomorphic to \mathbb{Z}_p . This implies $\check{\delta}_G = \theta^{d-1}$. \square

5.2 FINITELY GENERATED θ -ABELIAN PRO- p GROUPS

Recall that (G, θ) is said to be θ -abelian if $\ker(\theta) = Z_\theta(G)$ and $Z_\theta(G)$ is p -torsion free — in particular $\ker(\theta)$ is an abelian pro- p group. If G is finitely generated then one has an isomorphism of left $\mathbb{Z}_p[[G]]$ -modules $N \simeq \mathbb{Z}_p(1)^r$ for some non-negative integer r , and either $\Gamma = \mathrm{im}(\theta)$ is a finite group of order coprime to p , or Γ is a p -Poincaré duality group of dimension 1 satisfying $\check{\delta}_\Gamma = \mathbf{1}_\Gamma$ (cf. [23, Prop. 3.7.6]). Moreover, one has isomorphisms of left $\mathbb{Z}_p[[G]]$ -modules

$$H_k(N, \mathbb{Z}_p) \simeq \Lambda_k(N) \simeq \mathbb{Z}_p(k) \binom{r}{k}, \tag{5.3}$$

where $\Lambda_{\bullet}(_)$ denotes the exterior algebra over the ring \mathbb{Z}_p . Since $\text{cd}_p(\Gamma) \leq 1$, the Hochschild-Serre spectral sequence for homology (cf. [39, § 6.8])

$$E_{s,t}^2 = H_s(\Gamma, H_t(N, \mathbb{Z}_p(-m))) \implies H_{s+t}(G, \mathbb{Z}_p(-m)) \quad (5.4)$$

is concentrated in the first two columns. Hence, the spectral sequence collapses at the E^2 -term, i.e., $E_{s,t}^2 = E_{s,t}^\infty$. Thus, for $n \geq 1$ one has a short exact sequence

$$0 \longrightarrow H_{n-1}(N, \mathbb{Z}_p(-m))^\Gamma \longrightarrow H_n(G, \mathbb{Z}_p(-m)) \longrightarrow H_n(N, \mathbb{Z}_p(-m))_\Gamma \longrightarrow 0 \quad (5.5)$$

if $\text{cd}_p(\Gamma) = 1$, and isomorphisms

$$H^n(G, \mathbb{Z}_p(-m)) \simeq H_n(N, \mathbb{Z}_p(-m))_\Gamma \quad (5.6)$$

if Γ is a finite group of order coprime p . Here we used the fact that $H_0(\Gamma, _)= _-\Gamma$ coincides with the coinvariants of Γ , and that $H_1(\Gamma, _)= _-\Gamma$ coincides with the invariants of Γ if Γ is a p -Poincaré duality group of dimension 1 with $\delta_\Gamma = \mathbf{1}_\Gamma$. Since $H_{m-1}(N, \mathbb{Z}_p(-m))^\Gamma$ is a torsion free abelian pro- p group, and as

$$H_m(N, \mathbb{Z}_p(-m))_\Gamma = (H_m(N, \mathbb{Z}_p) \otimes \mathbb{Z}_p(-m))_\Gamma \simeq \Lambda_m(N) \quad (5.7)$$

by (5.3), one concludes from (5.5) and (5.6) that $H_m(G, \mathbb{Z}_p(-m))$ is torsion free.

PROPOSITION 5.3. *Let (G, θ) be a θ -abelian p -oriented virtual pro- p group such that $N = \ker(\theta)$ is a finitely generated torsion free abelian pro- p group, and that $\Gamma = \text{im}(\theta)$ is p -torsion free. Then G is a p -Poincaré duality group of dimension $d = \text{cd}(G)$, and θ is cyclotomic.*

Proof. By hypothesis, G is a p -torsion free p -adic analytic group. Hence the former assertion is a direct consequence of M. Lazard's theorem (cf. [33, Thm. 5.1.5]). The latter follows from Proposition 3.1. \square

From Proposition 5.2 one concludes the following:

COROLLARY 5.4. *Let (G, θ) be a θ -abelian pro- p group. If $p = 2$ assume further that $\text{im}(\theta)$ is torsion free.*

- (a) *The orientation θ is cyclotomic.*
- (b) *Suppose that G is finitely generated with minimum number of generators $d = d(G) < \infty$. If $p = 2$ assume further that $\text{im}(\theta) \subseteq 1 + 4\mathbb{Z}_2$. Then G is a Poincaré duality pro- p group of dimension d . Moreover, $\delta_G = \theta^{d-1}$.*
- (c) *If G satisfies the hypothesis of (b) and $d(G) \geq 2$, then for p odd, any cyclotomic orientation $\theta' : G \rightarrow \mathbb{Z}_p^\times$ of G must coincide with θ , i.e., $\theta' = \theta$. For $p = 2$ any cyclotomic orientation $\theta' : G \rightarrow \mathbb{Z}_2^\times$ satisfying $\text{im}(\theta') \subseteq 1 + 4\mathbb{Z}_2$ must coincide with θ .*

Proof. (a) follows from Proposition 5.3.

(b) By hypothesis, G is uniformly powerful (cf. [6, Ch. 4]), or equi- p -value, as it is called in [17]. Hence the claim follows from Proposition 5.3. By Proposition 5.2, $\bar{\delta}_G = \theta^{d-1}$.

(c) An element $\phi \in \text{Hom}_{\text{grp}}(G, \mathbb{Z}_p^\times)$ has finite order if, and only if, $\text{im}(\phi)$ is finite. Proposition 5.2 and part (b) imply that

$$\theta^{d-1} = \bar{\delta}_G = (\theta')^{d-1}.$$

Hence $(\theta^{-1}\theta')^{d-1} = \mathbf{1}_G$. For p odd, $\text{Hom}_{\text{grp}}(G, \mathbb{Z}_p^\times)$ does not contain non-trivial elements of finite order. Hence $\theta' = \theta$. For $p = 2$ the hypothesis implies that $\text{im}(\theta^{-1}\theta') \subseteq 1 + 4\mathbb{Z}_2$. Hence $(\theta^{-1}\theta')^{d-1} = \mathbf{1}_G$ implies that $\theta' = \theta$. \square

Note that, by Fact 2.2, Corollary 5.4(c) cannot hold if $d(G) = 1$.

5.3 PROFINITE p -POINCARÉ DUALITY GROUPS OF DIMENSION 2

As the following theorem shows, for a profinite p -Poincaré duality group G of dimension 2, the dualizing p -orientation $\bar{\delta}_G: G \rightarrow \mathbb{Z}_p^\times$ is always cyclotomic.

THEOREM 5.5. *Let G be a profinite p -Poincaré duality group of dimension 2. Then $\bar{\delta}_G: G \rightarrow \mathbb{Z}_p^\times$ is a cyclotomic p -orientation.*

Proof. As every p -oriented profinite group is 0-cyclotomic, it suffices to show that $H_{\text{cts}}^2(U, \mathbb{Z}_p(1))$ is torsion free for every open subgroup $U \subseteq G$. By Proposition 5.2, $\mathbb{Z}_p(-1) \simeq {}^\times D_G$. Hence, from the Eckmann-Shapiro lemma in homology and (5.1), one concludes that

$$\begin{aligned} H_1(U, \mathbb{Z}_p(-1)) &= \text{Tor}_1^U(\mathbb{Z}_p, \mathbb{Z}_p(-1)) \simeq \text{Tor}_1^U(\mathbb{Z}_p(-1)^\times, \mathbb{Z}_p) \\ &\simeq \text{Tor}_1^G(D_G, \mathbb{Z}_p[[G/U]]) \simeq H_{\text{cts}}^1(G, \mathbb{Z}_p[[G/U]]) \\ &\simeq \text{Hom}_{\text{grp}}(U, \mathbb{Z}_p). \end{aligned} \tag{5.8}$$

Hence $H_1(U, \mathbb{Z}_p(-1))$ is a torsion free \mathbb{Z}_p -module, and, by Proposition 3.1, $H_{\text{cts}}^2(U, \mathbb{Z}_p(1))$ is torsion free as well. \square

Remark 5.6. Let G be a profinite p -Poincaré duality group of dimension 2, and let $\bar{\delta}_G: G \rightarrow \mathbb{Z}_p^\times$ be the dualizing p -orientation. Then $(G, \bar{\delta}_G)$ is not necessarily Bloch-Kato, as the following example shows.

Let $p = 2$ and let $A = \text{PSL}_2(q)$ where $q \equiv 3 \pmod 4$. Then there exists a p -Frattini extension $\pi: G \rightarrow A$ of A such that G is a 2-Poincaré duality group of dimension 2, i.e., $\ker(\pi)$ is a pro-2 group contained in the Frattini subgroup of G (cf. [41]). In particular, G is perfect, and thus $\bar{\delta}_G = \mathbf{1}_G$. Hence $\mathbb{F}_2(1) = \mathbb{F}_2(0)$ is the trivial $\mathbb{F}_2[[G]]$ -module, and — as G is perfect — $H^1(G, \mathbb{F}_2(1)) = 0$. Moreover, $H^2(G, \mathbb{F}_2(2)) \simeq \mathbb{F}_2$, as G is a profinite 2-Poincaré duality group of dimension 2 with $\bar{\delta}_G = \mathbf{1}_G$. Therefore, $H^\bullet(G, \mathbf{1}_G)$ is not quadratic.

A pro- p group G which satisfies p -Poincaré duality in dimension 2 is also called a *Demuškin group* (cf. [23, Def. 3.9.9]). For this class of groups one has the following.

COROLLARY 5.7. *Let G be a Demuškin pro- p group. Then G is a Bloch-Kato pro- p group, and $\bar{\theta}_G: G \rightarrow \mathbb{Z}_p^\times$ is a cyclotomic p -orientation.*

Proof. By Theorem 5.5, it suffices to show that $(G, \bar{\theta}_G)$ is Bloch-Kato. It is well known that $H^\bullet(G, \hat{\theta}_G)$ is quadratic (cf. [31, §I.4.5]). Moreover, every open subgroup U of G is again a Demuškin group, with $\bar{\theta}_U = \bar{\theta}_G|_U$ (cf. [23, Thm. 3.9.15]). Hence $(G, \bar{\theta}_G)$ is Bloch-Kato. \square

Remark 5.8. [The Klein bottle pro-2 group] Let G be the pro-2 group given by the presentation

$$G = \langle x, y \mid xyx^{-1}y = 1 \rangle \quad (5.9)$$

Then G is a Demuškin pro-2 group containing the free abelian pro-2 group $H = \langle x^2, y \rangle$ of rank 2. Thus, by Corollary 5.7 $(G, \bar{\theta}_G)$ is cyclotomic. Since $H^1(G, \mathbb{I}_2(0)) \simeq \mathbb{I}_2 \oplus \mathbb{Z}/2\mathbb{Z}$, Proposition 3.1 implies that $\bar{\theta}_G \neq \mathbf{1}_G$ is non-trivial. In particular, since $\bar{\theta}_G|_H = \mathbf{1}_H$, this implies that $\text{im}(\bar{\theta}_G) = \{\pm 1\}$. Note that $H = \ker(\bar{\theta}_G)$ and that one has a canonical isomorphism

$$H = \langle x^2 \rangle \oplus \langle y \rangle \simeq \mathbb{Z}_2(0) \oplus \mathbb{Z}_2(1). \quad (5.10)$$

In particular, $(G, \bar{\theta}_G)$ is not $\bar{\theta}_G$ -abelian.

Example 5.9. Let G be the pro- p group with presentation

$$G = \langle x, y, z \mid [x, y] = z^{-p} \rangle.$$

If $p = 2$ then G is a Demuškin group, and $\bar{\theta}_G: G \rightarrow \mathbb{Z}_2^\times$ is given by $\bar{\theta}_G(x) = \bar{\theta}_G(y) = 1$, $\bar{\theta}_G(z) = -1$. On the other hand, if $p \neq 2$ then G is not a Demuškin group, and any p -orientations $\theta: G \rightarrow \mathbb{Z}_p^\times$ is not 1-cyclotomic (cf. [11, Thm. 8.1]). However, $H^\bullet(G, \hat{\theta})$ is still quadratic.

6 TORSION

It is well known that a Bloch-Kato pro- p group may have non-trivial torsion only if, $p = 2$. More precisely, a Bloch-Kato pro-2 group G is torsion if, and only if, G is abelian and of exponent 2. Moreover, any such group is a Bloch-Kato pro-2 group (cf. [25, §2]). The following result — which appeared first in [26, Prop. 2.13] — holds for 1-cyclotomically oriented pro- p groups (see also [11, Ex. 3.5] and [5, Ex. 14.27]).

PROPOSITION 6.1. *Let (G, θ) be a 1-cyclotomically oriented pro- p group.*

- (a) *If $\text{im}(\theta)$ is torsion free, then G is torsion free.*
- (b) *If G is non-trivial and torsion, then $p = 2$, $G \simeq C_2$ and θ is injective.*

Remark 6.2. Let $\theta: C_2 \rightarrow \mathbb{Z}_2^\times$ be an injective homomorphism of groups. Then $\mathbb{Z}_2(1) \simeq \omega_{C_2}$ is isomorphic to the augmentation ideal $\omega_{C_2} = \ker(\mathbb{Z}_2[C_2] \rightarrow \mathbb{Z}_2)$. Hence - by dimension shifting - $H^2(C_2, \mathbb{Z}_2(1)) = H^1(C_2, \mathbb{Z}_2(0)) = 0$. Thus - as C_2 has periodic cohomology of period 2 - one concludes that $H^s(C_2, \mathbb{Z}_2(t)) = 0$ for s odd and t even, and also for s even and t odd. Hence (C_2, θ) is cyclotomic. From Proposition 6.1 and the profinite version of Sylow's theorem one concludes the following corollary, which can be seen as a version of the Artin-Schreier theorem for 1-cyclotomically p -oriented profinite groups.

COROLLARY 6.3. *Let p be a prime number, and let (G, θ) be a profinite group with a 1-cyclotomic p -orientation.*

- (a) *If p is odd, then G has no p -torsion.*
- (b) *If $p = 2$, then every non-trivial 2-torsion subgroup is isomorphic to C_2 . Moreover, if $\text{im}(\theta)$ has no 2-torsion, then G has no 2-torsion.*

Remark 6.4. Let $\theta: \mathbb{Z}_2 \rightarrow \mathbb{Z}_2^\times$ be the homomorphism of groups given by $\theta(1 + \lambda) = -1$ and $\theta(\lambda) = 1$ for all $\lambda \in 2\mathbb{Z}_2$. Then θ is a 2-orientation of $G = \mathbb{Z}_2$ satisfying $\text{im}(\theta) = \{\pm 1\}$. As $\text{cd}_2(\mathbb{Z}_2) = 1$, Fact 2.2 implies that (\mathbb{Z}_2, θ) is Bloch-Kato and cyclotomically 2-oriented. However, $\text{im}(\theta)$ is not torsion free.

6.1 ORIENTATIONS ON $C_2 \times \mathbb{Z}_2$

As we have seen in Proposition 5.3, for p odd, every θ -abelian oriented pro- p group is cyclotomically p -oriented. For $p = 2$, this is not true. Indeed, one has the following.

PROPOSITION 6.5. *Any 2-orientation $\theta: G \rightarrow \mathbb{Z}_2^\times$ on $G \simeq C_2 \times \mathbb{Z}_2$ is not 1-cyclotomic.*

Proof. Suppose that (G, θ) is 1-cyclotomically 2-oriented. Let x, y be elements of G such that $x^2 = 1$ and $\text{ord}(y) = 2^\infty$, and that x, y generate G . Proposition 6.1 applied to the cyclic pro-2 group generated by x yields $\theta(x) = -1$. Put $\theta(y) = 1 + 2\lambda$ for some $\lambda \in \mathbb{Z}_2$. By [16, Prop. 6], if θ is 1-cyclotomic then for any pair of elements $c_x, c_y \in \mathbb{Z}_2(1)$ there exists a continuous crossed-homomorphism $c: G \rightarrow \mathbb{Z}_2(1)$ (i.e., a map satisfying $c(g_1 g_2) = c(g_1) + \theta(g_1)c(g_2)$, cf. [23, p. 15]) such that $c(x) = c_x$, $c(y) = c_y$. Set $c_x = c_y = 1$. Then one computes

$$\begin{aligned} c(xy) &= c_x + \theta(x)c_y = 1 - 1 = 0, & \text{and} \\ c(yx) &= c_y + \theta(y)c_x = 1 + 1 + 2\lambda, \end{aligned}$$

which yields $\lambda = -1$. The element xy has the same properties as y . Hence the previously mentioned argument applied to the element xy yields $\theta(xy) = 1 - 2 = -1$, whereas $\theta(xy) = \theta(x)\theta(y) = 1$, a contradiction. \square

Remark 6.6. From Proposition 6.1 and Proposition 6.5 one deduces that in a 1-cyclotomically 2-oriented pro-2 group, every element of order 2 is self-centralizing, which is a remarkable property of absolute Galois groups (cf. [4, Prop. 2.3] and [19, Cor. 2.3]).

PROPOSITION 6.7. *Let (G, θ) be a θ -abelian oriented pro-2 group. Then θ is cyclotomic if, and only if, either*

- (a) $\text{im}(\theta)$ is torsion free; or
- (b) $\text{im}(\theta)$ has order 2.

In both these cases (G, θ) is split θ -abelian.

Proof. Assume first that $\text{im}(\theta)$ is torsion free. Then the short exact sequence $\{1\} \rightarrow \ker(\theta) \rightarrow G \rightarrow \text{im}(\theta) \rightarrow \{1\}$ splits, as $\text{im}(\theta) \simeq \mathbb{Z}_2$ is a projective pro-2 group. Moreover, (G, θ) is cyclotomic by Proposition 5.3.

Second assume that θ is cyclotomic, $p = 2$ and that $\text{im}(\theta) \supseteq \{\pm 1\}$. If $g \in G$ satisfies $\theta(g) = -1$, then $g^2 \in \ker(\theta) = Z_\theta(G)$, and consequently

$$g^2 = g \cdot g^2 \cdot g^{-1} = (g^2)^{\theta(g)} = g^{-2},$$

i.e., $g^4 = 1$. Since $(\ker(\theta), \mathbf{1})$ is cyclotomically 2-oriented, $\ker(\theta)$ is torsion free, and one deduces that $g^2 = 1$. Therefore, the short exact sequence

$$\{1\} \longrightarrow H \longrightarrow G \longrightarrow C_2 \longrightarrow \{1\}$$

splits (here $H = \ker(\pi \circ \theta)$, where π is the canonical epimorphism $\mathbb{Z}_2^\times \twoheadrightarrow \{\pm 1\}$). Since $(H, \theta|_H)$ is again cyclotomically 2-oriented and as $\text{im}(\theta|_H)$ is torsion free, $(H, \theta|_H)$ is split $\theta|_H$ -abelian by the previously mentioned argument. We claim that $H = \ker(\theta)$. Indeed, suppose there exists $h \in H$ such that $\theta(h) \neq 1$. Put $\lambda = (1 + \theta(h))/2$ and let $z = ghgh^{-1} = [g, h^{-1}] \in \ker(\theta)$. Then - as $g = g^{-1}$ and $\theta(g) = -1$ - one has

$$\begin{aligned} g(z^\lambda h^2)g^{-1} &= (gzg)^\lambda \cdot gh^2g \\ &= z^{-\lambda} \cdot (ghg)^2 = z^{-\lambda} \cdot (ghgh^{-1} \cdot h)^2 \\ &= z^{-\lambda} \cdot (zhzh^{-1} \cdot h^2) = z^{-\lambda+1+\theta(h)} h^2 \\ &= z^\lambda h^2, \end{aligned}$$

i.e., g and $z^\lambda h^2$ commute which implies that $\langle g, z^\lambda h^2 \rangle \simeq C_2 \times \mathbb{Z}_p$ contradicting Proposition 6.5. Therefore, $H = \ker(\theta)$ is a free abelian pro-2 group, and $G \simeq H \rtimes C_2$.

Finally, let $p = 2$ and assume that $\text{im}(\theta) = \{\pm 1\}$. By Remark 6.2, we may also assume that $\ker(\theta)$ is non-trivial. Then, either

Case I: $\theta^{-1}(\{-1\})$ contains an element of order 2 and (G, θ) is split θ -abelian, i.e., $G \simeq \ker(\theta) \rtimes C_2$ with $\ker(\theta)$ a free abelian pro-2 group, or

Case II: all elements in $x \in \theta^{-1}(\{-1\})$ are of infinite order. Then for $y \in \ker(\theta)$, the group $K = \langle x, y \rangle$ must be isomorphic to the Klein bottle pro-2 group which is impossible as G is θ -abelian and thus contains only θ -abelian closed subgroups (cf. Remark 3.12(b)). Hence Case II is impossible.

By Lemma 3.10, if $U \subseteq G$ is an open subgroup, then either $U \subseteq \ker(\theta)$, or $U \simeq V \rtimes C_2$ for some open subgroup V of $\ker(\theta)$. In the first case, $(U, \mathbf{1})$ is

Remark 7.2. As before for a finitely generated pro- p group G let $d(G)$ denote its minimum number of generators. If p is odd and G is a finitely generated Bloch-Kato pro- p group, the cohomology ring $(H^\bullet(G, \mathbb{F}_p), \cup)$ is a quotient of the exterior \mathbb{F}_p -algebra $\Lambda_\bullet = \Lambda_\bullet(H^1(G, \mathbb{F}_p))$. In particular, $\text{cd}_p(G) \leq d(G)$. Moreover, $\Lambda_{d(G)}$ is the unique minimal ideal of Λ_\bullet . Hence equality of $\text{cd}_p(G)$ and $d(G)$ is equivalent to $H^\bullet(G, \mathbb{F}_p)$ being isomorphic to Λ_\bullet . It is well known that this implies that G is uniformly powerful (cf. [33, Thm. 5.1.6]), and that there exists a p -orientation $\theta: G \rightarrow \mathbb{Z}_p^\times$ such that G is θ -abelian (cf. [25, Thm. 4.6]). Let $p = 2$, and let (G, θ) be a cyclotomically oriented Bloch-Kato pro-2 group satisfying $\text{im}(\theta) \subseteq 1 + 4\mathbb{Z}_2$. Then Proposition 7.1 implies that the cohomology ring $(H^\bullet(G, \mathbb{F}_2), \cup)$ is a quotient of the exterior \mathbb{F}_2 -algebra $\Lambda_\bullet = \Lambda_\bullet(H^1(G, \mathbb{F}_2))$, and hence $\text{cd}_2(G) \leq d(G)$. If $\text{cd}_2(G) = d(G)$, the previously mentioned argument, Proposition 7.1 and [42] imply that G is uniformly powerful. Finally, [25, Thm. 4.11] yields that G is θ' -abelian for some orientation $\theta': G \rightarrow \mathbb{Z}_2^\times$. Thus, if $d(G) \geq 2$, one has $\theta = \theta'$ by Corollary 5.4(c).

From the above remark and J-P. Serre's theorem (cf. [30]) one concludes the following fact.

FACT 7.3. *Let (G, θ) be a finitely generated cyclotomically oriented torsion free Bloch-Kato pro-2 group. Then $\text{cd}_2(G) < \infty$.*

7.1 TITS' ALTERNATIVE

From Remark 7.2 one concludes the following.

FACT 7.4. (a) *Let p be odd, and let G be a Bloch-Kato pro- p group satisfying $d(G) \leq 2$. Then G is either isomorphic to a free pro- p group, or G is θ -abelian for some orientation $\theta: G \rightarrow \mathbb{Z}_p^\times$.*

(b) *Let $p = 2$, and let (G, θ) be a cyclotomically oriented Bloch-Kato pro-2 group satisfying $\text{im}(\theta) \subseteq 1 + 4\mathbb{Z}_2$ and $d(G) \leq 2$. Then G is either isomorphic to a free pro-2 group, or G is θ -abelian.*

In [25, Thm. 4.6] it was shown, that for p odd any Bloch-Kato pro- p group satisfies a strong form of Tits' alternative (cf. [35]), i.e., either G contains a closed non-abelian free pro- p subgroup, or there exists a p -orientation $\theta: G \rightarrow \mathbb{Z}_p^\times$ such that G is θ -abelian. Using the results from the previous subsection and [25, Thm. 4.11], one obtains the following version of Tits' alternative if p is equal to 2.

PROPOSITION 7.5. *Let (G, θ) be a cyclotomically oriented virtual pro-2 group which is also Bloch-Kato, such that $\text{im}(\theta) \subseteq 1 + 4\mathbb{Z}_2$. Then either G contains a closed non-abelian free pro-2 subgroup; or G is θ -abelian.*

Proof. As $\text{im}(\theta) \subseteq 1 + 4\mathbb{Z}_2$, Proposition 6.1-(a) implies that G is torsion free. From Proposition 7.1 one concludes that the first Bockstein morphism β^1 vanishes. Thus, the hypothesis of [25, Thm. 4.11] are satisfied (cf. Remark 7.2), and this yields the claim. \square

Remark 7.6. Note that Proposition 7.5 without the hypothesis $\text{im}(\theta) \subseteq 1 + 4\mathbb{Z}_2$ does not remain true (cf. Remark 5.8).

7.2 THE θ -CENTER

One has the following characterization of the θ -center for a cyclotomically oriented Bloch-Kato pro- p group (G, θ) .

THEOREM 7.7. *Let (G, θ) be a cyclotomically oriented torsion free Bloch-Kato pro- p group. If $p = 2$ assume further that $\text{im}(\theta) \subseteq 1 + 4\mathbb{Z}_2$. Then $Z_\theta(G)$ is the unique maximal closed abelian normal subgroup of G contained in $\ker(\theta)$.*

Proof. Let $A \subseteq \ker(\theta)$ be a closed abelian normal subgroup of G , let $z \in A$, $z \neq 1$, and let $x \in G$ be an arbitrary element. Put $C = \text{cl}(\langle x, z \rangle) \subseteq G$. Then either $C \simeq \mathbb{Z}_p$ or C is a 2-generated pro- p group. Thus, by Fact 7.4, one has to distinguish three cases:

- (i) $d(C) = 1$;
- (ii) $d(C) = 2$ and C is isomorphic to a free pro- p group; or
- (iii) $d(C) = 2$ and C is θ' -abelian for some p -orientation $\theta': C \rightarrow \mathbb{Z}_p^\times$.

In case (i), x and z commute. If C is generated by z , then $C \subseteq \ker(\theta)$ and $\theta(x) = 1$. If C is generated by x , then $z = x^\lambda$ for some $\lambda \in \mathbb{Z}_p$, and $1 = \theta(z) = \theta(x)^\lambda$. Hence $\theta(x) = 1$, as $\text{im}(\theta)$ is torsion free. In both cases $xzx^{-1} = z = z^{\theta(x)}$.

Case (ii) cannot hold: by hypothesis, $A \cap C \neq \{1\}$, but free pro- p groups of rank 2 do not contain non-trivial closed abelian normal subgroups.

Suppose that case (iii) holds. Then $\theta' = \theta|_C$ by Corollary 5.4(c), and $z \in \ker(\theta|_C) = Z_{\theta|_C}(C)$. Therefore, $xzx^{-1} = z^{\theta|_C(x)} = z^{\theta(x)}$.

Hence we have shown that for all $z \in A$ and all $x \in G$ one has that $xzx^{-1} = z^{\theta(x)}$. This yields the claim. \square

The above result can be seen as the group theoretic generalization of [12, Corollary 3.3] and [13, Thm. 4.6]. Note that in the case $p = 2$ the additional hypothesis in Theorem 7.7 is necessary (cf. Remark 5.8). Indeed, if G is the Klein bottle pro-2 group then $\langle x^2 \rangle$ is another maximal closed abelian normal subgroup of G contained in $\ker(\partial_G)$.

Remark 7.8. Let \mathbb{K} be a field containing a primitive p^{th} -root of unity. Theorem 7.7, together with [12, Thm. 3.1] and [13, Thm. 4.6], implies that the $\theta_{\mathbb{K}, p}$ -center of the maximal pro- p Galois group $G_{\mathbb{K}}(p)$ is the inertia group of the maximal p -henselian valuation admitted by \mathbb{K} .

7.3 ISOLATED SUBGROUPS

Let G be a pro- p group, and let $S \subseteq G$ be a closed subgroup of G . Then S is called *isolated*, if for all $g \in G$ for which there exists $k \geq 1$ such that $g^{p^k} \in S$ follows that $g \in S$. Hence a closed normal subgroup N of G is isolated if, and only if, G/N is torsion free.

PROPOSITION 7.9. *Let (G, θ) be an oriented Bloch-Kato pro- p group. In the case $p = 2$ assume further that $\text{im}(\theta) \subseteq 1 + 4\mathbb{Z}_2$ and that θ is 1-cyclotomic. Then $Z_\theta(G)$ is an isolated subgroup of G .*

Proof. Suppose there exists $x \in G \setminus Z_\theta(G)$ and $k \geq 1$ such that $x^{p^k} \in Z_\theta(G)$. By changing the element x if necessary, we may assume that $k = 1$, i.e., $x^p \in Z_\theta(G)$. As G is torsion free (cf. Corollary 6.3), one has that $x^p \neq 1$.

For an arbitrary $g \in G$, the subgroup $C(g) = \text{cl}(\langle g, x \rangle) \subseteq G$ is not free, as $gx^pg^{-1} = x^{p\theta(g)}$. Thus, from Fact 7.4 one concludes that $C(g)$ is $\theta|_{C(g)}$ -abelian. Moreover, as $\text{im}(\theta)$ is torsion-free, $\theta(x^p) = \theta(x)^p = 1$ implies that $x \in \ker(\theta|_{C(g)}) = Z_{\theta|_{C(g)}}(C(g))$. Thus, $x \in \bigcap_{g \in G} Z_{\theta|_{C(g)}}(C(g)) \subseteq Z_\theta(G)$. \square

Proposition 7.9 generalises to profinite groups as follows.

COROLLARY 7.10. *Let (G, θ) be a torsion free p -oriented Bloch-Kato profinite group. For $p = 2$ assume also that $\text{im}(\theta) \subseteq 1 + 4\mathbb{Z}_2$ and that θ is 1-cyclotomic. Then $Z_\theta(G)$ is an isolated subgroup of G .*

Proof. Let $x \in Z_\theta(G)$, $y \in G$ and $n \in \mathbb{N}$ such that $x = y^n$. Then $Y = \text{cl}(\langle y \rangle)$ is pro-cyclic and virtually pro- p . Thus, as G is torsion free by hypothesis, Y is a cyclic pro- p group, and n is a p -power. Let $P \in \text{Syl}_p(G)$ be a pro- p Sylow subgroup of G containing Y . Then $(P, \theta|_P)$ satisfies the hypothesis of Proposition 7.9, which yields the claim. \square

7.4 SPLIT EXTENSIONS

PROPOSITION 7.11. *Let (G, θ) be a p -oriented Bloch-Kato pro- p group of finite cohomological dimension satisfying $\text{im}(\theta) \subseteq 1 + p\mathbb{Z}_p$ (resp. $\text{im}(\theta) \subseteq 1 + 4\mathbb{Z}_2$ if $p = 2$), and let Z be a closed normal subgroup of G isomorphic to \mathbb{Z}_p such that G/Z is torsion free. Then $Z \not\subseteq G^p[G, G]$.*

Proof. Let $d = \text{cd}_p(G)$. As $\text{cd}(Z) = 1$, and as $H^1(Z, \mathbb{F}_p) \simeq \mathbb{F}_p$, one has $\text{vcd}_p(G/Z) = d - 1$ (cf. [43]). Thus, as G/Z is torsion free, J-P. Serre's theorem (cf. [30]) implies that $\text{cd}_p(G/Z) = d - 1$.

Suppose that $Z \subseteq G^p[G, G]$. Then $\text{inf}_{G,Z}^1: H^1(G/Z, \mathbb{F}_p) \rightarrow H^1(G, \mathbb{F}_p)$ is an isomorphism. For $\chi \in H^1(G, \mathbb{F}_p)$, set $\bar{\chi} \in H^1(G/Z, \mathbb{F}_p)$ such that $\chi = \text{inf}_{G,Z}^1(\bar{\chi})$. Then, by [23, Prop. 1.5.3] one has

$$\chi_1 \cup \dots \cup \chi_k = \text{inf}_{G,Z}^1(\bar{\chi}_1) \cup \dots \cup \text{inf}_{G,Z}^1(\bar{\chi}_k) = \text{inf}_{G,Z}^k(\bar{\chi}_1 \cup \dots \cup \bar{\chi}_k)$$

for any $\chi_1, \dots, \chi_k \in H^1(G, \mathbb{F}_p)$, i.e.,

$$\text{inf}_{G,Z}^k: H^k(G/Z, \mathbb{F}_p) \longrightarrow H^k(G, \mathbb{F}_p) \tag{7.2}$$

is surjective for all $k \geq 0$. Let

$$(E_r^{st}, d_r) \Rightarrow H^{s+t}(G, \mathbb{F}_p), \quad E_2^{st} = H^s(G/Z, H^t(Z, \mathbb{F}_p)) \tag{7.3}$$

denote the Hochschild-Serre spectral sequence associated to the extension of pro- p groups $Z \rightarrow G \rightarrow G/Z$ with coefficients in the discrete G -module \mathbb{F}_p . We claim that E_∞^{st} is concentrated on the bottom row, i.e., $E_\infty^{st} = 0$ for all $t \geq 1$. Since $\text{cd}_p(Z) = 1$ and $\text{cd}_p(G/Z) = d - 1$, one has $E_2^{st} = 0$ for $t \geq 2$ or $s \geq d$. Hence, d_r^{st} is the 0-map for every $s, t \geq 0$ and $r \geq 3$, i.e., $E_\infty^{st} \simeq E_3^{st}$. The total complex $\text{tot}_\bullet(E_\infty^{\bullet\bullet})$ of the graded \mathbb{F}_p -bialgebra $E_\infty^{\bullet\bullet}$ coincides with $H^\bullet(G, \mathbb{F}_p)$, which is quadratic by hypothesis. Thus $E_\infty^{\bullet\bullet}$ is generated by

$$\text{tot}_1(E_\infty^{\bullet\bullet}) = E_\infty^{1,0} = E_2^{1,0}.$$

Hence, $E_3^{st} = 0$ for $t \geq 1$.

On the other hand, $H^1(Z, \mathbb{F}_p)$ is a trivial G/Z -module isomorphic to \mathbb{F}_p , and thus, as $\text{cd}_p(G/Z) = d - 1$, one has

$$E_2^{d-1,1} = H^{d-1}(G/Z, H^1(Z, \mathbb{F}_p)) \neq 0. \tag{7.4}$$

Moreover, $d_2^{d-1,1}$ is the 0-map, thus $E_3^{d-1,1} = \ker(d_2^{d-1,1}) = E_\infty^{d-1,1} \neq 0$, a contradiction, and this yields the claim. \square

Proposition 7.11 has the following consequence.

PROPOSITION 7.12. *Let (G, θ) be a p -oriented Bloch-Kato pro- p group (resp. virtual pro- p group) of finite cohomological p -dimension, and let Z be a closed normal subgroup of G isomorphic to \mathbb{Z}_p such that G/Z is torsion free. Then there exists a Z -complement C in G , i.e., the extension of profinite groups*

$$\{1\} \longrightarrow Z \longrightarrow G \longrightarrow G/Z \longrightarrow \{1\} \tag{7.5}$$

splits.

Proof. Assume first that G is a pro- p group. By Proposition 7.11, one has that $Z \not\subseteq \Phi(G) = G^p[G, G]$. Hence there exists a maximal closed subgroup C_1 of G such that $C_1 Z = G$ and $Z_1 = C_1 \cap Z = Z^p$. Moreover, Z_1 is a closed normal subgroup in C_1 such that C_1/Z_1 is torsion free and $Z_1 \simeq \mathbb{Z}_p$. From Proposition 7.11 again, one concludes that $Z_1 \not\subseteq \Phi(C_1)$. Thus repeating this process one finds open subgroup C_k of G of index p^k such that $C_k Z = G$ and $Z_k = C_k \cap Z = Z^{p^k}$. Hence $C = \bigcap_{k \geq 1} C_k$ is a Z -complement in G .

If G is a p -oriented virtual pro- p group, then G is a $\bar{\Sigma}$ -virtual pro- p group for $\bar{\Sigma} = \text{im}(\theta)$ (cf. 4.1), and thus corresponds to $(O_p(G), \theta^\circ, \gamma)$ in alternative form. In particular, the maximal subgroup C_1 and hence all closed subgroups C_k can be chosen to be $\bar{\Sigma}$ -invariant (cf. Proposition 4.5). Hence $C = \bigcap_{k \in \mathbb{N}} C_k$ carries canonically a left $\bar{\Sigma}$ -action, and thus defines a Z complement $H = C \rtimes \bar{\Sigma}$ in G . \square

The proof of Theorem 1.2 can be deduced from Proposition 7.12 as follows.

Proof of Theorem 1.2. Assume first that G is either pro- p , or virtually pro- p . To prove statement (i) (and (ii)), we proceed by induction on $d = \text{cd}_p(G) = \text{cd}(G)$. For $d = 1$, G is free (resp. virtually free) (cf. [23, Prop. 3.5.17]), and thus $Z_\theta(G) = \{1\}$. So assume that $d \geq 1$, and that the claim holds for $d - 1$. Note that $Z_\theta(G)$ is a finitely generated abelian pro- p group satisfying $d_\circ = d(Z_\theta(G)) = \text{cd}_p(Z_\theta(G)) \leq d$. If $d_\circ = 0$, there is nothing to prove. If $d_\circ \geq 1$, $Z_\theta(G)$ contains an isolated closed subgroup Z satisfying $d(Z) = 1$. By definition, Z is normal in G . Hence Proposition 7.12 implies that there exists a subgroup $C \subseteq G$ satisfying $C \cap Z = \{1\}$ and $CZ = G$. As $C \simeq G/Z$, the main result of [43] implies that $\text{cd}(C) = \text{vcd}(C) = d - 1$. Since $Z_{\theta|_C}(C)Z = Z_\theta(G)$, the claim then follows by induction.

To prove statement (iii), let $G^\circ = \ker(\hat{\theta}: G \rightarrow \mathbb{F}_p^\times)$ and $\bar{G}^\circ = \ker(\hat{\theta}: \bar{G} \rightarrow \mathbb{F}_p^\times)$, and put $\bar{O} = O^p(\bar{G}^\circ)$ and

$$O = \{g \in G^\circ \mid gZ_\theta(G) \in \bar{O}^p(\bar{G})\}. \tag{7.6}$$

Then, by construction, $\text{im}(\hat{\theta}|_{\bar{O}})$ is a pro- p group and hence trivial. In particular, the left $\mathbb{F}_p[[\bar{O}]]$ -module $\mathbb{F}_p(1)$ is the trivial module. Thus, as \bar{O} is p -perfect, one concludes that

$$H^1(\bar{O}, \mathbb{F}_p(1)) = 0. \tag{7.7}$$

By hypothesis, $(\bar{G}, \hat{\theta})$ is Bloch-Kato, and therefore $(\bar{O}, \mathbf{1})$ is Bloch-Kato. Hence (7.7) yields that

$$H^k(\bar{O}, \mathbb{F}_p(j)) = H^k(\bar{O}, \mathbb{F}_p(0)) = 0 \tag{7.8}$$

for all positive integers k, j . Note that $\mathbb{Z}_p(1)$ is the trivial $\mathbb{Z}_p[[\bar{O}]]$ -module isomorphic to \mathbb{Z}_p as abelian pro- p group. The cyclotomicity of $(\bar{O}, \mathbf{1})$ implies that $H^2(\bar{O}, \mathbb{Z}_p(1))$ is p -torsion free, and from the exact sequence

$$0 \longrightarrow H^2(\bar{O}, \mathbb{Z}_p(1)) \xrightarrow{p} H^2(\bar{O}, \mathbb{Z}_p(1)) \longrightarrow H^2(\bar{O}, \mathbb{F}_p(1)) \longrightarrow 0 \tag{7.9}$$

one concludes that

$$H^2(\bar{O}, \mathbb{Z}_p(1)) = 0. \tag{7.10}$$

By hypothesis, $\text{cd}_p(Z_\theta(G)) \leq \text{cd}_p(G) < \infty$, and thus $Z_\theta(G) \simeq \mathbb{Z}_p(1)^r$ is a trivial left $\mathbb{Z}_p[[\bar{O}]]$ -module and a finitely generated free (abelian pro- p group). Hence

$$H^2(\bar{O}, Z_\theta(G)) = 0, \tag{7.11}$$

which implies that

$$\{1\} \longrightarrow Z_\theta(G) \longrightarrow O \xrightarrow{\pi} \bar{O} \longrightarrow \{1\} \tag{7.12}$$

is a split short exact sequence of profinite groups. From this fact one concludes that

$$O = Z_\theta(G) \cdot O^p(G^\circ) \quad \text{and} \quad Z_\theta(G) \cap O^p(G^\circ) = \{1\}. \tag{7.13}$$

Let $\tilde{G} = G/O^p(G^\circ)$. Then for all abelian pro- p groups M with a continuous left $\mathbb{Z}_p[[\tilde{G}]]$ -action inflation induces an isomorphism in cohomology

$$\text{inf}_{\tilde{G}}^G(-): H_{\text{cts}}^k(\tilde{G}, M) \longrightarrow H_{\text{cts}}^k(G, M) \tag{7.14}$$

(cf. Proposition 4.6). Moreover, as $\theta|_O = \mathbf{1}$ is the constant 1 function, θ induces a p -orientation $\tilde{\theta}: \tilde{G} \rightarrow \mathbb{Z}_p^\times$ on \tilde{G} . In particular, from (7.14) one concludes that $\text{cd}_p(\tilde{G}) < \infty$, and that $(\tilde{G}, \tilde{\theta})$ is cyclotomic and Bloch-Kato. Thus, by part (i), the exact sequence of virtual pro- p groups

$$\{1\} \longrightarrow Z_\theta(G)O^p(G^\circ)/O^p(G^\circ) \longrightarrow \tilde{G} \xrightarrow{\tilde{\pi}} \tilde{G}/\bar{O} \longrightarrow \{1\} \tag{7.15}$$

splits. Let $\tilde{H} \subset \tilde{G}$ be a complement for $Z_\theta(G)O^p(G^\circ)/O^p(G^\circ)$ in \tilde{G} , and let

$$H = \{g \in G^\circ \mid gO^p(G^\circ) \in \tilde{H}\}. \tag{7.16}$$

Then, by construction, $H \cap Z_\theta(G)O^p(G^\circ) \subseteq O^p(G^\circ)$. Thus $HO^p(G^\circ)$ is a complement of $Z_\theta(G)$ in G . □

Finally, we ask whether the converse of Theorem 3.13 holds true.

QUESTION 7.13. *Let (G, θ) be a cyclotomically p -oriented Bloch-Kato pro- p group, and suppose that*

$$H^\bullet(G, \mathbb{F}_p) \simeq H^\bullet(C, \mathbb{F}_p) \otimes \Lambda_\bullet(V),$$

for some subgroup $C \subseteq G$ and some nontrivial subspace $V \subseteq H^1(G, \mathbb{F}_p)$. Does there exist an isolated closed subgroup $Z \subseteq Z_\theta(G)$ such that $G = CZ$ and $Z/Z^p \simeq V^ = \text{Hom}(V, \mathbb{F}_p)$?*

7.5 THE ELEMENTARY TYPE CONJECTURE

In order to formulate a conjecture concerning the maximal pro- p Galois groups of fields, I. Efrat introduced in [9] the class \mathcal{C}_{FG} of p -oriented pro- p groups (resp. cyclotomic pro- p pairs) of *elementary type*.

This class consists of all finitely generated p -oriented pro- p groups which can be constructed from \mathbb{Z}_p and Demuškin groups using coproducts and fibre products (cf. [9, § 3]).

Efrat’s *elementary type conjecture* asks whether every pair $(G_{\mathbb{K}}(p), \theta_{\mathbb{K}, p})$ for which \mathbb{K} contains a primitive p^{th} -root of unity and $G_{\mathbb{K}}(p)$ is finitely generated, belongs to \mathcal{C}_{FG} (see [7], and also [15] for the case $p = 2$). This conjecture originates from the theory of quadratic forms (cf. [20], [10, p. 268]).

One may extend slightly Efrat’s class by defining the class \mathcal{E}_{CO} of *cyclotomically p -oriented Bloch-Kato pro- p groups of elementary type* to be the smallest class of cyclotomically p -oriented pro- p groups containing

- (a) (F, θ) , with F a finitely generated free pro- p group and $\theta: F \rightarrow \mathbb{Z}_p^\times$ any p -orientation;

- (b) $(G, \bar{\delta}_G)$, with G a Demuškin pro- p group;
- (c) $(\mathbb{Z}/2\mathbb{Z}, \theta)$, with $\text{im}(\theta) = \{\pm 1\}$ in case that $p = 2$;

and which is closed under coproducts and under fibre products with respect to finitely generated split θ -abelian pro- p groups, i.e., if (G_1, θ_1) and (G_2, θ_2) are contained in \mathcal{E}_{CO} , then

- (d) $(G, \theta) = (G_1, \theta_1) \amalg (G_2, \theta_2) \in \mathcal{E}_{\text{CO}}$; and
- (e) $(G, \theta) = \mathbb{Z}_p \rtimes_{\theta_1} (G_1, \theta_1) \in \mathcal{E}_{\text{CO}}$.

Question 1.5 asks whether every finitely generated cyclotomically p -oriented Bloch-Kato pro- p group belongs to the class \mathcal{E}_{CO} . By Theorem 1.1, Question 1.5 is stronger than Efrat's elementary type conjecture. Nevertheless, it is stated in purely group theoretic terms.

Remark 7.14. Recently, Question 1.5 has received a positive solution in the class of *trivially p -oriented right-angled Artin pro- p groups*: I. Snopce and P.A. Zalesskiĭ proved that the only indecomposable right-angled Artin pro- p group which is Bloch-Kato and cyclotomically p -oriented is $(\mathbb{Z}_p, \mathbf{1})$ (cf. [32]).

ACKNOWLEDGEMENTS

The authors are grateful to: the anonymous referee, for her/his valuable suggestions; to I. Efrat, for the interesting discussion they had together at the Ben-Gurion University of the Negev in 2016; and to D. Neftin and I. Snopce, for their interest. Also, the first-named author wishes to thank M. Florence and P. Guillot for the discussions on the preprint [5].

Both authors were partially supported by the PRIN 2015 "Group Theory and Applications". The first-named author was also partially supported by the Israel Science Foundation (grant No. 152/13).

REFERENCES

- [1] S. Bloch and K. Kato. p -adic étale cohomology. *Inst. Hautes Études Sci. Publ. Math.*, 63:107–152, 1986.
- [2] K. S. Brown. *Cohomology of groups*, volume 87 of *Graduate Texts in Mathematics*. Springer-Verlag, New York-Berlin, 1982.
- [3] A. Brumer. Pseudocompact algebras, profinite groups and class formations. *J. Algebra*, 4:442–470, 1966.
- [4] T. C. Craven and T. L. Smith. Formally real fields from a Galois-theoretic perspective. *J. Pure Appl. Algebra*, 145(1):19–36, 2000.
- [5] C. De Clercq and M. Florence. Lifting theorems and smooth profinite groups. preprint, available at [arxiv:1710.10631](https://arxiv.org/abs/1710.10631), 2017.

- [6] J. D. Dixon, M. P. F. du Sautoy, A. Mann, and D. Segal. *Analytic pro- p groups*, volume 61 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 1999.
- [7] I. Efrat. Orderings, valuations, and free products of Galois groups. *Sem. Structure Algébriques Ordonnées, Univ. Paris VII*, 1995.
- [8] I. Efrat. Pro- p Galois groups of algebraic extensions of \mathbf{Q} . *J. Number Theory*, 64(1):84–99, 1997.
- [9] I. Efrat. Small maximal pro- p Galois groups. *Manuscripta Math.*, 95(2):237–249, 1998.
- [10] I. Efrat. *Valuations, orderings, and Milnor K -theory*, volume 124 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2006.
- [11] I. Efrat and C. Quadrelli. The Kummerian property and maximal pro- p Galois groups. *J. Algebra*, 525:284–310, 2019.
- [12] A. J. Engler and J. Koenigsmann. Abelian subgroups of pro- p Galois groups. *Trans. Amer. Math. Soc.*, 350(6):2473–2485, 1998.
- [13] A. J. Engler and J. B. Nogueira. Maximal abelian normal subgroups of Galois pro-2-groups. *J. Algebra*, 166(3):481–505, 1994.
- [14] M. Fried and M. Jarden. *Field arithmetic*, volume 11 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, third edition, 2008. Revised by Jarden.
- [15] B. Jacob and R. Ware. A recursive description of the maximal pro-2 Galois group via Witt rings. *Math. Z.*, 200(3):379–396, 1989.
- [16] J. P. Labute. Classification of Demushkin groups. *Canad. J. Math.*, 19:106–132, 1967.
- [17] M. Lazard. Groupes analytiques p -adiques. *Inst. Hautes Études Sci. Publ. Math.*, 26:389–603, 1965.
- [18] S. Mac Lane. *Homology*. Classics in Mathematics. Springer-Verlag, Berlin, 1995. Reprint of the 1975 edition.
- [19] L. Mahé, J. Mináč, and T. L. Smith. Additive structure of multiplicative subgroups of fields and Galois theory. *Doc. Math.*, 9:301–355, 2004.
- [20] M. Marshall. The elementary type conjecture in quadratic form theory. In *Algebraic and arithmetic theory of quadratic forms*, volume 344 of *Contemp. Math.*, pages 275–293. Amer. Math. Soc., Providence, RI, 2004.

- [21] J. Mináč and N. D. Tân. Triple Massey products vanish over all fields. *J. Lond. Math. Soc. (2)*, 94(3):909–932, 2016.
- [22] J. Mináč and N. D. Tân. Triple Massey products and Galois theory. *J. Eur. Math. Soc. (JEMS)*, 19(1):255–284, 2017.
- [23] J. Neukirch, A. Schmidt, and K. Wingberg. *Cohomology of number fields*, volume 323 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 2008.
- [24] A. Polishchuk and L. Positselski. *Quadratic algebras*, volume 37 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2005.
- [25] C. Quadrelli. Bloch-Kato pro- p groups and locally powerful groups. *Forum Math.*, 26(3):793–814, 2014.
- [26] C. Quadrelli. *Cohomology of Absolute Galois Groups*. PhD thesis, University of Western Ontario, 2015.
- [27] L. Ribes and P. A. Zalesskiĭ. *Profinite groups*, volume 40. Springer-Verlag, Berlin, 2000.
- [28] J-E. Roos. Sur les foncteurs dérivés de \varprojlim . Applications. *C. R. Acad. Sci. Paris*, 252:3702–3704, 1961.
- [29] M. Rost. Norm varieties and algebraic cobordism. In *Proceedings of the International Congress of Mathematicians. Vol. II (Beijing, 2002)*, pages 77–85. Higher Ed. Press, Beijing, 2002.
- [30] J.-P. Serre. Sur la dimension cohomologique des groupes profinis. *Topology*, 3:413–420, 1965.
- [31] J.-P. Serre. *Galois cohomology*. Springer-Verlag, Berlin, 1997.
- [32] I. Snopce and P. A. Zalesskiĭ. Right-angled artin pro- p groups. preprint, available at [arXiv:2005.01685](https://arxiv.org/abs/2005.01685), 2020.
- [33] P. Symonds and T. Weigel. Cohomology of p -adic analytic groups. In *New horizons in pro- p groups*, volume 184 of *Progr. Math.*, pages 349–410. Birkhäuser Boston, Boston, MA, 2000.
- [34] J. Tate. Relations between K_2 and Galois cohomology. *Invent. Math.*, 36:257–274, 1976.
- [35] J. Tits. Free subgroups in linear groups. *J. Algebra*, 20:250–270, 1972.
- [36] V. Voevodsky. On motivic cohomology with \mathbf{Z}/l -coefficients. *Ann. of Math. (2)*, 174(1):401–438, 2011.

- [37] A. R. Wadsworth. p -Henselian field: K -theory, Galois cohomology, and graded Witt rings. *Pacific J. Math.*, 105(2):473–496, 1983.
- [38] R. Ware. Galois groups of maximal p -extensions. *Trans. Amer. Math. Soc.*, 333(2):721–728, 1992.
- [39] C. A. Weibel. *An introduction to homological algebra*, volume 38 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1994.
- [40] C. A. Weibel. The norm residue isomorphism theorem. *J. Topol.*, 2(2):346–372, 2009.
- [41] T. Weigel. Frattini extensions and class field theory. In *Groups St. Andrews 2005. Vol. 2*, volume 340 of *London Math. Soc. Lecture Note Ser.*, pages 661–684. Cambridge Univ. Press, Cambridge, 2007.
- [42] T. Weigel. A characterization of powerful pro-2 groups by their cohomology. preprint, 2016.
- [43] T. Weigel and P. A. Zalesskiĭ. Profinite groups of finite cohomological dimension. *C. R., Math., Acad. Sci. Paris*, 338(5):353–358, 2004.

Claudio Quadrelli
Department of Mathematics and
Applications
Università di Milano-Bicocca
Via R.Cozzi 55 - ed. U5
20125 Milan
Italy
claudio.quadrelli@unimib.it

Thomas S. Weigel
Department of Mathematics and
Applications
Università di Milano-Bicocca
Via R.Cozzi 55 - ed. U5
20125 Milan
Italy
thomas.weigel@unimib.it